

## t22\_nfcont\_4

(TML3g7vz5K6CMh3XT4iVBR5ixqbvJShN92Z)

October 27, 2020

Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $v1\_xreal\_0 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_euclid : \iota \Rightarrow \iota$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_nfcont\_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $r2\_nfcont\_4 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k7\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_fcont\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k4\_real\_ns1 : \iota \Rightarrow \iota$  be given. Let  $r2\_nfcont\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v2\_struct\_0 : \iota \Rightarrow o$  be given. Let  $v13\_algstr\_0 : \iota \Rightarrow o$  be given. Let  $v2\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v4\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v5\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v6\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v7\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v8\_rlvect\_1 : \iota \Rightarrow o$  be given. Let  $v3\_normsp\_0 : \iota \Rightarrow o$  be given. Let  $v4\_normsp\_0 : \iota \Rightarrow o$  be given. Let  $v2\_normsp\_1 : \iota \Rightarrow o$  be given. Let  $l1\_normsp\_1 : \iota \Rightarrow o$  be given. Let  $r1\_nfcont\_3 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $v1\_normsp\_1 : \iota \Rightarrow o$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $k5\_euclid : \iota \Rightarrow \iota$  be given. Let  $r1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $u1\_algstr\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_real\_ns1 : \iota \Rightarrow \iota$  be given. Let  $u1\_rlvect\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_real\_ns1 : \iota \Rightarrow \iota$  be given. Let  $u1\_normsp\_0 : \iota \Rightarrow \iota$  be given. Let  $k3\_real\_ns1 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(m2\_subset\_1 X0 k1\_numbers k5\_numbers) \Rightarrow (\forall X1. \\
 & ((v1\_funct\_1 X1) \wedge (m1\_subset\_1 X1 (k1\_zfmisc\_1 (k2\_zfmisc\_1 ( \\
 & \quad k1\_euclid X0) k1\_numbers)))) \Rightarrow (\forall X2. ((v1\_funct\_1 X2) \wedge ( \\
 & m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 (u1\_struct\_0 (k4\_real\_ns1 \\
 & \quad X0)) k1\_numbers)))) \Rightarrow (\forall X3. (m1\_subset\_1 X3 (k1\_euclid X0)) \Rightarrow \\
 & (\forall X4. (m1\_subset\_1 X4 (u1\_struct\_0 (k4\_real\_ns1 X0))) \Rightarrow \\
 & (((X1 = X2) \wedge (X3 = X4)) \Rightarrow ((r2\_nfcont\_4 X0 X1 X3) \Leftrightarrow (r2\_nfcont\_1 (k4\_real\_ns1 \\
 & \quad X0) X2 X4))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
& \forall X0.(v1\_xreal\_0 X0) \Rightarrow (\forall X1.((\neg v2\_struct\_0 X1) \wedge \\
& (v13\_algstr\_0 X1) \wedge ((v2\_rlvect\_1 X1) \wedge ((v3\_rlvect\_1 X1) \wedge ((v4\_rlvect\_1 \\
& X1) \wedge ((v5\_rlvect\_1 X1) \wedge ((v6\_rlvect\_1 X1) \wedge ((v7\_rlvect\_1 X1) \wedge \\
& ((v8\_rlvect\_1 X1) \wedge ((v3\_normsp\_0 X1) \wedge ((v4\_normsp\_0 X1) \wedge ((v2\_normsp\_1 \\
& X1) \wedge (l1\_normsp\_1 X1)))))))))) \Rightarrow (\forall X2.((v1\_funct\_1 \\
& X2) \wedge (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k1\_numbers (u1\_struct\_0 \\
& X1)))))) \Rightarrow (\forall X3.((v1\_funct\_1 X3) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 \\
& (k2\_zfmisc\_1 (u1\_struct\_0 X1) k1\_numbers)))) \Rightarrow (((X0 \in k1\_relset\_1 \\
& k1\_numbers (k1\_partfun1 k1\_numbers (u1\_struct\_0 X1) (u1\_struct\_0 \\
& X1) k1\_numbers X2 X3)) \wedge ((r1\_nfcont\_3 X1 X2 X0) \wedge (r2\_nfcont\_1 X1 \\
& X3 (k7\_partfun1 (u1\_struct\_0 X1) X2 X0)))) \Rightarrow (r1\_fcont\_1 (k1\_partfun1 \\
& k1\_numbers (u1\_struct\_0 X1) (u1\_struct\_0 X1) k1\_numbers X2 X3 \\
& X0))))))
\end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0.(m2\_subset\_1 X0 k1\_numbers k5\_numbers) \Rightarrow (\forall X1. \\
& (v1\_xreal\_0 X1) \Rightarrow (\forall X2.((v1\_funct\_1 X2) \wedge (m1\_subset\_1 X2 \\
& (k1\_zfmisc\_1 (k2\_zfmisc\_1 k1\_numbers (k1\_euclid X0)))))) \Rightarrow (\forall X3. \\
& ((v1\_funct\_1 X3) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 k1\_numbers \\
& (u1\_struct\_0 (k4\_real\_ns1 X0)))))) \Rightarrow ((X3 = X2) \Rightarrow ((r1\_nfcont\_4 \\
& X0 X2 X1) \Leftrightarrow (r1\_nfcont\_3 (k4\_real\_ns1 X0) X3 X1))))))
\end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned}
& \forall X0.\forall X1.((\neg v1\_xboole\_0 X0) \wedge ((\neg v1\_xboole\_0 X1) \wedge \\
& (m1\_subset\_1 X1 (k1\_zfmisc\_1 X0)))) \Rightarrow (\forall X2.(m2\_subset\_1 \\
& X2 X0 X1) \Leftrightarrow (m1\_subset\_1 X2 X1))
\end{aligned} \tag{4}$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \tag{5}$$

Assume the following.

$$(\neg v1\_xboole\_0 k4\_ordinal1) \wedge (v3\_ordinal1 k4\_ordinal1) \tag{6}$$

Assume the following.

$$\begin{aligned}
& \forall X0.(v7\_ordinal1 X0) \Rightarrow ((\neg v2\_struct\_0 (k4\_real\_ns1 X0)) \wedge \\
& ((v13\_algstr\_0 (k4\_real\_ns1 X0)) \wedge ((v2\_rlvect\_1 (k4\_real\_ns1 \\
& X0)) \wedge ((v3\_rlvect\_1 (k4\_real\_ns1 X0)) \wedge ((v4\_rlvect\_1 (k4\_real\_ns1 \\
& X0)) \wedge ((v5\_rlvect\_1 (k4\_real\_ns1 X0)) \wedge ((v6\_rlvect\_1 (k4\_real\_ns1 \\
& X0)) \wedge ((v7\_rlvect\_1 (k4\_real\_ns1 X0)) \wedge ((v8\_rlvect\_1 (k4\_real\_ns1 \\
& X0)) \wedge ((v3\_normsp\_0 (k4\_real\_ns1 X0)) \wedge ((v4\_normsp\_0 (k4\_real\_ns1 \\
& X0)) \wedge ((v1\_normsp\_1 (k4\_real\_ns1 X0)) \wedge (v2\_normsp\_1 (k4\_real\_ns1 \\
& X0))))))))))
\end{aligned} \tag{7}$$

Assume the following.

$$\neg v1\_xboole\_0 \ k1\_numbers \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1\_relat\_1 \ X1)\wedge((v5\_relat\_1 \ X1 \ X0)\wedge(v1\_funct\_1 \ X1)))\Rightarrow(m1\_subset\_1 \ (k7\_partfun1 \ X0 \ X1 \ X2) \ X0) \quad (9)$$

Assume the following.

$$m1\_subset\_1 \ k5\_numbers \ (k1\_zfmisc\_1 \ k1\_numbers) \quad (10)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 \ X0)\Rightarrow((\neg v2\_struct\_0 \ (k4\_real\_ns1 \ X0))\wedge((v1\_normsp\_1 \ (k4\_real\_ns1 \ X0))\wedge(l1\_normsp\_1 \ (k4\_real\_ns1 \ X0)))) \quad (11)$$

Assume the following.

$$\begin{aligned} \forall X0.(v7\_ordinal1 \ X0)\Rightarrow(\forall X1.((\neg v2\_struct\_0 \ X1)\wedge \\ ((v1\_normsp\_1 \ X1)\wedge(l1\_normsp\_1 \ X1)))\Rightarrow((X1 = k4\_real\_ns1 \ X0)\Leftrightarrow \\ ((u1\_struct\_0 \ X1 = k1\_euclid \ X0)\wedge((k4\_struct\_0 \ X1 = k5\_euclid \ X0)\wedge \\ ((r1\_funct\_2 \ (k2\_zfmisc\_1 \ (u1\_struct\_0 \ X1) \ (u1\_struct\_0 \ X1)) \\ (u1\_struct\_0 \ X1) \ (k2\_zfmisc\_1 \ (k1\_euclid \ X0) \ (k1\_euclid \ X0)) \ ( \\ k1\_euclid \ X0) \ (u1\_algstr\_0 \ X1) \ (k1\_real\_ns1 \ X0))\wedge((r1\_funct\_2 \\ (k2\_zfmisc\_1 \ k1\_numbers \ (u1\_struct\_0 \ X1)) \ (u1\_struct\_0 \ X1) \ (k2\_zfmisc\_1 \\ k1\_numbers \ (k1\_euclid \ X0)) \ (k1\_euclid \ X0) \ (u1\_rlvect\_1 \ X1) \ (k2\_real\_ns1 \\ X0))\wedge(r1\_funct\_2 \ (u1\_struct\_0 \ X1) \ k1\_numbers \ (k1\_euclid \ X0) \ k1\_numbers \\ (u1\_normsp\_0 \ X1) \ (k3\_real\_ns1 \ X0))))))))) \quad (12) \end{aligned}$$

Assume the following.

$$\forall X0.(m1\_subset\_1 \ X0 \ k4\_ordinal1)\Rightarrow(v7\_ordinal1 \ X0) \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 \ X2 \ (k1\_zfmisc\_1 \ (k2\_zfmisc\_1 \ X0 \ X1)))\Rightarrow((v4\_relat\_1 \ X2 \ X0)\wedge(v5\_relat\_1 \ X2 \ X1)) \quad (14)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 \ X2 \ (k1\_zfmisc\_1 \ (k2\_zfmisc\_1 \ X0 \ X1)))\Rightarrow(v1\_relat\_1 \ X2) \quad (15)$$

### Theorem 1

$$\begin{aligned} \forall X0.(m2\_subset\_1 \ X0 \ k1\_numbers \ k5\_numbers)\Rightarrow(\forall X1. \\ (v1\_xreal\_0 \ X1)\Rightarrow(\forall X2.((v1\_funct\_1 \ X2)\wedge(m1\_subset\_1 \ X2 \\ (k1\_zfmisc\_1 \ (k2\_zfmisc\_1 \ k1\_numbers \ (k1\_euclid \ X0))))))\Rightarrow(\forall X3. \\ ((v1\_funct\_1 \ X3)\wedge(m1\_subset\_1 \ X3 \ (k1\_zfmisc\_1 \ (k2\_zfmisc\_1 \ ( \\ k1\_euclid \ X0) \ k1\_numbers))))\Rightarrow(((X1 \in k1\_relset\_1 \ k1\_numbers \ ( \\ k1\_partfun1 \ k1\_numbers \ (k1\_euclid \ X0) \ (k1\_euclid \ X0) \ k1\_numbers \\ X2 \ X3))\wedge((r1\_nfcont\_4 \ X0 \ X2 \ X1)\wedge(r2\_nfcont\_4 \ X0 \ X3 \ (k7\_partfun1 \\ (k1\_euclid \ X0) \ X2 \ X1))))\Rightarrow(r1\_fcont\_1 \ (k1\_partfun1 \ k1\_numbers \\ (k1\_euclid \ X0) \ (k1\_euclid \ X0) \ k1\_numbers \ X2 \ X3) \ X1)))))) \end{aligned}$$