

t22_qc_lang2
(TMJ3T5HaCyWd591YmEUMrja3EzdUWi1BdLV)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k3_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k8_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k15_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_qc_lang2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m2_subset_1 X1 (k2_qc_lang1 \\ X0) (k3_qc_lang1 X0)) \Rightarrow (\forall X2.(m2_subset_1 X2 (k2_qc_lang1 \\ X0) (k3_qc_lang1 X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 (k9_qc_lang1 \\ X0)) \Rightarrow (\forall X4.(m1_subset_1 X4 (k9_qc_lang1 X0)) \Rightarrow ((k15_qc_lang1 \\ X0 X1 X3 = k15_qc_lang1 X0 X2 X4) \Rightarrow ((X1 = X2) \wedge (X3 = X4))))))))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (\neg v1_xboole_0 (k3_qc_lang1 X0)) \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.\forall X3.((m1_qc_lang1 X0) \wedge \\ ((m1_subset_1 X1 (k3_qc_lang1 X0)) \wedge ((m1_subset_1 X2 (k3_qc_lang1 \\ X0)) \wedge (m1_subset_1 X3 (k9_qc_lang1 X0)))))) \Rightarrow (m1_subset_1 (k6_qc_lang2 \\ X0 X1 X2 X3) (k9_qc_lang1 X0)) \end{aligned} \quad (4)$$

Assume the following.

$$\forall X0.(m1_qc_lang1 X0) \Rightarrow (m1_subset_1 (k3_qc_lang1 X0) (k1_zfmisc_1 \\ (k2_qc_lang1 X0))) \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k2_qc_lang1 \\ & X0)\ (k3_qc_lang1\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k2_qc_lang1 \\ & X0)\ (k3_qc_lang1\ X0)) \Rightarrow (\forall X3.(m2_subset_1\ X3\ (k2_qc_lang1 \\ & X0)\ (k3_qc_lang1\ X0)) \Rightarrow (\forall X4.(m1_subset_1\ X4\ (k9_qc_lang1 \\ & X0)) \Rightarrow (k8_qc_lang2\ X0\ X1\ X2\ X3\ X4 = k15_qc_lang1\ X0\ X1\ (k6_qc_lang2 \\ & X0\ X2\ X3\ X4)))))) \end{aligned} \quad (6)$$

Assume the following.

$$\forall X0.(v1_xboole_0\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ X0)) \Rightarrow (v1_xboole_0\ X1)) \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_qc_lang1\ X0) \Rightarrow (\forall X1.(m2_subset_1\ X1\ (k2_qc_lang1 \\ & X0)\ (k3_qc_lang1\ X0)) \Rightarrow (\forall X2.(m2_subset_1\ X2\ (k2_qc_lang1 \\ & X0)\ (k3_qc_lang1\ X0)) \Rightarrow (\forall X3.(m2_subset_1\ X3\ (k2_qc_lang1 \\ & X0)\ (k3_qc_lang1\ X0)) \Rightarrow (\forall X4.(m1_subset_1\ X4\ (k9_qc_lang1 \\ & X0)) \Rightarrow (\forall X5.(m1_subset_1\ X5\ (k9_qc_lang1\ X0)) \Rightarrow (\forall X6. \\ & (m2_subset_1\ X6\ (k2_qc_lang1\ X0)\ (k3_qc_lang1\ X0)) \Rightarrow ((k8_qc_lang2 \\ & X0\ X1\ X2\ X3\ X4 = k15_qc_lang1\ X0\ X6\ X5) \Rightarrow ((X1 = X6) \wedge (k6_qc_lang2\ X0\ X2 \\ & X3\ X4 = X5)))))))))) \end{aligned}$$