

t22_qc_lang3 (TMZvXzRtzorPXLXRCHfNwTh- MXby9h17kYKF)

October 27, 2020

Let $m1_qc_lang1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $v6_qc_lang1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k14_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k24_qc_lang1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_qc_lang1 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.k2_xboole_0 X0 k1_xboole_0 = X0 \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(k2_xboole_0 X0 X1 = k1_xboole_0) \Rightarrow (X0 = k1_xboole_0) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(m1_qc_lang1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k9_qc_lang1 \\ X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (k9_qc_lang1 X0)) \Rightarrow (k24_qc_lang1 \\ X0 (k14_qc_lang1 X0 X1 X2) = k4_subset_1 (k3_qc_lang1 X0) (k24_qc_lang1 \\ X0 X1) (k24_qc_lang1 X0 X2)))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((m1_subset_1 X1 (k1_zfmisc_1 \\ X0)) \wedge (m1_subset_1 X2 (k1_zfmisc_1 X0))) \Rightarrow (k4_subset_1 X0 X1 X2 = \\ k2_xboole_0 X1 X2) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.((m1_qc_lang1 X0) \wedge (m1_subset_1 X1 (k9_qc_lang1 \\ X0))) \Rightarrow (m1_subset_1 (k24_qc_lang1 X0 X1) (k1_zfmisc_1 (k3_qc_lang1 \\ X0))) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((m1_qc_lang1\ X0)\wedge((m1_subset_1\ X1\ (k9_qc_lang1\ X0))\wedge(m1_subset_1\ X2\ (k9_qc_lang1\ X0))))\Rightarrow(m1_subset_1\ (k14_qc_lang1\ X0\ X1\ X2)\ (k9_qc_lang1\ X0)) \quad (6)$$

Assume the following.

$$\forall X0.(m1_qc_lang1\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ (k9_qc_lang1\ X0))\Rightarrow((v6_qc_lang1\ X1\ X0)\Leftrightarrow(k24_qc_lang1\ X0\ X1 = k1_xboole_0))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.k2_xboole_0\ X0\ X1 = k2_xboole_0\ X1\ X0 \quad (8)$$

Theorem 1

$$\forall X0.(m1_qc_lang1\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ (k9_qc_lang1\ X0))\Rightarrow(\forall X2.(m1_subset_1\ X2\ (k9_qc_lang1\ X0))\Rightarrow(((v6_qc_lang1\ X1\ X0)\wedge(v6_qc_lang1\ X2\ X0))\Leftrightarrow(v6_qc_lang1\ (k14_qc_lang1\ X0\ X1\ X2)\ X0))))$$