

t22_sprect_3 (TMFm-
LAM1Qj1ZPAyxu8zcfWGz9xkoedqnMA1)

October 27, 2020

Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_topreal1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_jordan3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_1 : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_jordan3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m1_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $v6_membered : \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v1_card_1 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & ((X1 \in k3_topreal1 np_2 X0) \Rightarrow ((r1_xxreal_0 np_1 (k1_jordan3 X0 \\ & X1)) \wedge (\neg r1_xxreal_0 (k3_finseq_1 X0) (k1_jordan3 X0 X1)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 X1) \wedge ((\neg v2_xxreal_0 X0) \wedge (\neg v3_xxreal_0 X1)))) \quad (2)$$

Assume the following.

$$\forall X0.((\neg v1_xboole_0 X0) \wedge ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0)))) \Rightarrow ((np_1 \in k4_finseq_1 X0) \wedge (k3_finseq_1 X0 \in k4_finseq_1 X0)) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(r1_xxreal_0 X0 X1) \wedge (\neg v2_xxreal_0 X1) \wedge (v2_xxreal_0 X0))) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & ((X1 \in k3_topreal1 np_2 X0) \Rightarrow (((X1 \neq k1_funct_1 X0 np_1) \Rightarrow (k3_finseq_1 \\ & (k3_jordan3 X0 X1) = k2_nat_1 (k1_jordan3 X0 X1) np_1)) \wedge ((X1 = k1_funct_1 \\ & X0 np_1) \Rightarrow (k3_finseq_1 (k3_jordan3 X0 X1) = k1_jordan3 X0 X1)))))) \quad (5) \end{aligned}$$

Assume the following.

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & ((X1 \in k3_topreal1 np_2 X0) \Rightarrow ((k1_funct_1 (k3_jordan3 X0 X1) (k3_finseq_1 \\ & (k3_jordan3 X0 X1)) = X1) \wedge (\forall X2.(m1_subset_1 X2 k5_numbers) \Rightarrow \\ & (((r1_xxreal_0 np_1 X2) \wedge (r1_xxreal_0 X2 (k1_jordan3 X0 X1))) \Rightarrow \\ & (k1_funct_1 (k3_jordan3 X0 X1) X2 = k1_funct_1 X0 X2)))))) \quad (6) \end{aligned}$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.\forall X2.(m2_finseq_1 X2 X1) \Rightarrow (((r1_xxreal_0 np_1 X0) \wedge (r1_xxreal_0 X0 (k3_finseq_1 X2))) \Rightarrow (k7_partfun1 X1 X2 X0 = k1_funct_1 X2 X0))) \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (r1_xxreal_0 X0 (k2_xcmplx_0 X0 X1))) \quad (8)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (9)$$

Assume the following.

$$r1_xxreal_0 np_1 np_1 \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1 X1 X0) \Leftrightarrow (m1_finseq_1 X1 X0) \quad (11)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (12)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (13)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k3_finseq_1 X0 = k1_card_1 X0) \quad (14)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_subset_1 X0 k5_numbers) \wedge (v7_ordinal1 X1)) \Rightarrow (k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (15)$$

Assume the following.

$$v6_membered k4_ordinal1 \quad (16)$$

Assume the following.

$$v3_membered k1_numbers \quad (17)$$

Assume the following.

$$\forall X0. (v1_xboole_0 X0) \Rightarrow ((v1_xboole_0 (k1_card_1 X0)) \wedge (v1_card_1 (k1_card_1 X0))) \quad (18)$$

Assume the following.

$$\forall X0. \forall X1. (m2_finseq_1 X1 X0) \Rightarrow ((v1_funct_1 X1) \wedge ((v1_finseq_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers X0)))))) \quad (19)$$

Assume the following.

$$\forall X0. \forall X1. (m1_finseq_1 X1 X0) \Rightarrow ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1))) \quad (20)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))) \wedge (m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (m2_finseq_1 (k3_jordan3 X0 X1) (u1_struct_0 (k15_euclid np_2))) \quad (21)$$

Assume the following.

$$\forall X0. \forall X1. ((m1_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))) \wedge (m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2)))) \Rightarrow (m1_subset_1 (k1_jordan3 X0 X1) k5_numbers) \quad (22)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge((v5_relat_1 X1 X0)\wedge(v1_funct_1 X1)))\Rightarrow(\forall X2.(X2 \in k9_xtuple_0 X1)\Rightarrow(k7_partfun1 X0 X1 X2 = k1_funct_1 X1 X2)) \quad (23)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0)\wedge(v1_xcmplx_0 X1))\Rightarrow(k2_xcmplx_0 X0 X1 = k2_xcmplx_0 X1 X0) \quad (24)$$

Assume the following.

$$\forall X0.((v3_ordinal1 X0)\wedge(v1_finset_1 X0))\Rightarrow(v7_ordinal1 X0) \quad (25)$$

Assume the following.

$$\forall X0.(v3_membered X0)\Rightarrow(v1_membered X0) \quad (26)$$

Assume the following.

$$\forall X0.((v1_xxreal_0 X0)\wedge(v2_xxreal_0 X0))\Rightarrow((\neg v1_xboole_0 X0)\wedge((v1_xxreal_0 X0)\wedge(\neg v3_xxreal_0 X0))) \quad (27)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xcmplx_0 X0) \quad (28)$$

Assume the following.

$$\forall X0.\forall X1.(m1_finseq_1 X1 X0)\Rightarrow(v5_relat_1 X1 X0) \quad (29)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_xxreal_0 X0) \quad (30)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_xreal_0 X0) \quad (31)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0)\Rightarrow(v1_finset_1 X0) \quad (32)$$

Assume the following.

$$\forall X0.(v1_card_1 X0)\Rightarrow(v3_ordinal1 X0) \quad (33)$$

Assume the following.

$$\forall X0.(v6_membered X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(v7_ordinal1 X1)) \quad (34)$$

Assume the following.

$$\forall X0.(v1_membered X0)\Rightarrow(\forall X1.(m1_subset_1 X1 X0)\Rightarrow(v1_xcmplx_0 X1)) \quad (35)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m2_finseq_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & ((X1 \in k3_topreal1 np_2 X0) \Rightarrow (k7_partfun1 (u1_struct_0 (k15_euclid \\ & np_2)) (k3_jordan3 X0 X1) np_1 = k7_partfun1 (u1_struct_0 (k15_euclid \\ & np_2)) X0 np_1))) \end{aligned}$$