

t22_tmap_1
(TMKynXp89tvon1BRY4L5i5ijdRw59rtE8pg)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $m1_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_tsep_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k2_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_pre_topc : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. \\ & (m1_pre_topc X1 X0) \Rightarrow (\forall X2.(m1_pre_topc X2 X0) \Rightarrow ((r1_tarski \\ & (u1_struct_0 X1) (u1_struct_0 X2)) \Leftrightarrow (m1_pre_topc X1 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((r1_tarski X0 X1) \wedge \\ & (r1_tarski X2 X3)) \Rightarrow (r1_tarski (k2_xboole_0 X0 X2) (k2_xboole_0 \\ & X1 X3)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(((\neg v2_struct_0 X0) \wedge (l1_pre_topc \\ & X0)) \wedge (((\neg v2_struct_0 X1) \wedge (m1_pre_topc X1 X0)) \wedge ((\neg v2_struct_0 \\ & X2) \wedge (m1_pre_topc X2 X0)))) \Rightarrow ((\neg v2_struct_0 (k1_tsep_1 X0 X1 X2)) \wedge \\ & ((v1_pre_topc (k1_tsep_1 X0 X1 X2)) \wedge (m1_pre_topc (k1_tsep_1 X0 \\ & X1 X2) X0))) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_pre_topc X0)) \Rightarrow (\forall X1. \\ & ((\neg v2_struct_0 X1) \wedge (m1_pre_topc X1 X0)) \Rightarrow (\forall X2.((\neg v2_struct_0 \\ & X2) \wedge (m1_pre_topc X2 X0)) \Rightarrow (\forall X3.((\neg v2_struct_0 X3) \wedge ((v1_pre_topc \\ & X3) \wedge (m1_pre_topc X3 X0)) \Rightarrow ((X3 = k1_tsep_1 X0 X1 X2) \Leftrightarrow (u1_struct_0 \\ & X3 = k2_xboole_0 (u1_struct_0 X1) (u1_struct_0 X2)))))) \end{aligned} \quad (4)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v2_pre_topc X0) \wedge (l1_pre_topc \\ & X0))) \Rightarrow (\forall X1.((\neg v2_struct_0 X1) \wedge (m1_pre_topc X1 X0)) \Rightarrow (\\ & \forall X2.((\neg v2_struct_0 X2) \wedge (m1_pre_topc X2 X0)) \Rightarrow (\forall X3. \\ & ((\neg v2_struct_0 X3) \wedge (m1_pre_topc X3 X0)) \Rightarrow (\forall X4.((\neg v2_struct_0 \\ & X4) \wedge (m1_pre_topc X4 X0)) \Rightarrow (((m1_pre_topc X1 X2) \wedge (m1_pre_topc \\ & X3 X4)) \Rightarrow (m1_pre_topc (k1_tsep_1 X0 X1 X3) (k1_tsep_1 X0 X2 X4)))))) \end{aligned}$$