

t22\_xxreal\_0  
(TMF1ViVycQTrbd3snQHaj9Hmg4iBGwb11Db)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow ((r1\_xxreal\_0 X0 X1) \wedge (r1\_xxreal\_0 X1 X2)) \Rightarrow \\ & (r1\_xxreal\_0 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (r1\_xxreal\_0 (k3\_xxreal\_0 X0 X1) X0)) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow (v1\_xxreal\_0 (k3\_xxreal\_0 X0 X1)) \quad (3)$$

**Theorem 1**

$$\begin{aligned} & \forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (\forall X2. \\ & (v1\_xxreal\_0 X2) \Rightarrow ((r1\_xxreal\_0 X0 (k3\_xxreal\_0 X1 X2)) \Rightarrow (r1\_xxreal\_0 \\ & X0 X1)))) \end{aligned}$$