

t22_xxreal_1

(TMZU4XBGrjfrxmkenFVJHygymTTEccdK8vJ)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_xxreal_0 X1) \Rightarrow (\forall X2. (v1_xxreal_0 X2) \Rightarrow ((X0 \in k4_xxreal_1 X1 X2) \Rightarrow ((X0 \in k2_xxreal_1 X1 X2) \wedge (X0 \neq X1)))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (2)$$

Theorem 1

$$\forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow (r1_tarski (k4_xxreal_1 X0 X1) (k2_xxreal_1 X0 X1)))$$