

t230_xreal_1 (TM-
 MVsM8CvT1JN18v3dRkKQcRv3nJ9Nz11QL)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 k6_numbers X0) \wedge (r1_xxreal_0 \\ & X1 X2)) \Rightarrow (r1_xxreal_0 X1 (k2_xcmplx_0 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 X0 (k2_xcmplx_0 X1 X2)) \Rightarrow (r1_xxreal_0 \\ & (k6_xcmplx_0 X0 X1) X2)))) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (((r1_xxreal_0 k6_numbers X0) \wedge (r1_xxreal_0 \\ & X1 X2)) \Rightarrow (r1_xxreal_0 (k6_xcmplx_0 X1 X0) X2)))) \end{aligned}$$