

t232\_xcplx\_1

(TMHyXso4wbg2kVHk6T5pgJrwEmvrzfsDaBH)

October 27, 2020

Let  $v1\_xcplx\_0 : \iota \Rightarrow o$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k7\_xcplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xcplx\_0 X0) \Rightarrow (\forall X1.(v1\_xcplx\_0 X1) \Rightarrow ((X0 \neq k6\_numbers) \Rightarrow (k7\_xcplx\_0 X0 (k7\_xcplx\_0 X0 X1) = X1))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xcplx\_0 X0) \wedge (v1\_xcplx\_0 X1)) \Rightarrow (v1\_xcplx\_0 (k7\_xcplx\_0 X0 X1)) \quad (2)$$

**Theorem 1**

$$\forall X0.(v1\_xcplx\_0 X0) \Rightarrow (\forall X1.(v1\_xcplx\_0 X1) \Rightarrow (\neg (X1 \neq k6\_numbers) \wedge (\forall X2.(v1\_xcplx\_0 X2) \Rightarrow (X0 \neq k7\_xcplx\_0 X1 X2))))$$