

t23_circrm1
(TMGu8a7xmyQLusYaWEvjCQ8FFTvLxQ1n67z)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_msualg_1 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $r1_circrm1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u4_struct_0 : \iota \Rightarrow \iota$ be given. Let $v2_funct_1 : \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k2_funct_1 : \iota \Rightarrow \iota$ be given. Let $r3_pua2mss1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u2_msualg_1 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $u1_msualg_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v2_funct_1 X0) \Rightarrow \\ ((k10_xtuple_0 X0 = k9_xtuple_0 (k2_funct_1 X0)) \wedge (k9_xtuple_0 \\ X0 = k10_xtuple_0 (k2_funct_1 X0)))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. (v1_relat_1 X0) \Rightarrow (k7_relat_1 X0 (k9_xtuple_0 X0) = k10_xtuple_0 X0) \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v1_relat_1 (k2_funct_1 X0)) \wedge (v1_funct_1 (k2_funct_1 X0))) \quad (3)$$

Assume the following.

$$\begin{aligned} \forall X0. ((\neg v2_struct_0 X0) \wedge (l1_msualg_1 X0)) \Rightarrow (\forall X1. \\ ((\neg v2_struct_0 X1) \wedge (l1_msualg_1 X1)) \Rightarrow (\forall X2. ((v1_relat_1 \\ X2) \wedge (v1_funct_1 X2)) \Rightarrow (\forall X3. ((v1_relat_1 X3) \wedge (v1_funct_1 \\ X3)) \Rightarrow ((r1_circrm1 X0 X1 X2 X3) \Leftrightarrow ((v2_funct_1 X2) \wedge ((v2_funct_1 \\ X3) \wedge (r3_pua2mss1 X0 X1 X2 X3) \wedge (r3_pua2mss1 X1 X0 (k2_funct_1 X2) \\ (k2_funct_1 X3)))))))))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned}
& \forall X0.(l1_msualg_1 X0) \Rightarrow (\forall X1.(l1_msualg_1 X1) \Rightarrow (\forall X2. \\
& ((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow (\forall X3.((v1_relat_1 \\
& X3) \wedge (v1_funct_1 X3)) \Rightarrow ((r3_pua2mss1 X0 X1 X2 X3) \Leftrightarrow ((k9_xtuple_0 \\
& X2 = u1_struct_0 X0) \wedge ((k9_xtuple_0 X3 = u4_struct_0 X0) \wedge ((r1_tarski \\
& (k10_xtuple_0 X2) (u1_struct_0 X1)) \wedge ((r1_tarski (k10_xtuple_0 \\
& X3) (u4_struct_0 X1)) \wedge ((k3_relat_1 (u2_msualg_1 X0) X2 = k3_relat_1 \\
& X3 (u2_msualg_1 X1)) \wedge (\forall X4. \forall X5. ((v1_relat_1 X5) \wedge \\
& (v1_funct_1 X5)) \Rightarrow (((X4 \in u4_struct_0 X0) \wedge (X5 = k1_funct_1 (u1_msualg_1 \\
& X0) X4)) \Rightarrow (k3_relat_1 X5 X2 = k1_funct_1 (u1_msualg_1 X1) (k1_funct_1 \\
& X3 X4))))))))))))) \\
& \tag{5}
\end{aligned}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge (l1_msualg_1 X0)) \Rightarrow (\forall X1. \\
& ((\neg v2_struct_0 X1) \wedge (l1_msualg_1 X1)) \Rightarrow (\forall X2.((v1_relat_1 \\
& X2) \wedge (v1_funct_1 X2)) \Rightarrow (\forall X3.((v1_relat_1 X3) \wedge (v1_funct_1 \\
& X3)) \Rightarrow ((r1_cictrm1 X0 X1 X2 X3) \Rightarrow ((u1_struct_0 X1 = k7_relat_1 X2 \\
& (u1_struct_0 X0)) \wedge (u4_struct_0 X1 = k7_relat_1 X3 (u4_struct_0 \\
& X0)))))))))
\end{aligned}$$