

t23_facirc_1

(TMQqBp2hArPNWphsriAKz643VGET1S7wAoA)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v1_circcomb : \iota \Rightarrow o$ be given. Let $v2_circcomb : \iota \Rightarrow o$ be given. Let $l1_msualg_1 : \iota \Rightarrow o$ be given. Let $k2_circcomb : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_circcomb : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge (l1_msualg_1 X0)) \Rightarrow (\forall X1. \\ & ((\neg v2_struct_0 X1) \wedge (l1_msualg_1 X1)) \Rightarrow ((r1_circcomb X0 X1) \Rightarrow (\\ & k2_circcomb X0 X1 = k2_circcomb X1 X0))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_circcomb X0) \wedge ((v2_circcomb \\ & X0) \wedge (l1_msualg_1 X0)))) \Rightarrow (\forall X1.((\neg v2_struct_0 X1) \wedge ((v1_circcomb \\ & X1) \wedge ((v2_circcomb X1) \wedge (l1_msualg_1 X1)))) \Rightarrow (r1_circcomb X0 X1)) \end{aligned} \quad (2)$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v1_circcomb X0) \wedge ((v2_circcomb \\ & X0) \wedge (l1_msualg_1 X0)))) \Rightarrow (\forall X1.((\neg v2_struct_0 X1) \wedge ((v1_circcomb \\ & X1) \wedge ((v2_circcomb X1) \wedge (l1_msualg_1 X1)))) \Rightarrow (k2_circcomb X0 X1 = \\ & k2_circcomb X1 X0)) \end{aligned}$$