

t23_funct_7

(TMM3qgsG45tikCeFQGF9h1CQqSE3CzS1hBi)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v1_funcop_1 : \iota \Rightarrow o$ be given. Let $k7_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k4_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. ((X0 \in X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X2))) \Rightarrow (m1_subset_1 X0 X2) \quad (1)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (2)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (k4_finseq_1 X0 = k9_xtuple_0 X0) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_funcop_1 X0))) \Rightarrow ((v1_relat_1 (k1_funct_1 X0 X1)) \wedge (v1_funct_1 (k1_funct_1 X0 X1))) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \wedge ((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 X1)))) \Rightarrow ((v1_relat_1 (k7_finseq_1 X0 X1)) \wedge ((v1_funct_1 (k7_finseq_1 X0 X1)) \wedge (v1_finseq_1 (k7_finseq_1 X0 X1)))) \quad (5)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow (m1_subset_1 (k4_finseq_1 X0) (k1_zfmisc_1 k5_numbers)) \quad (6)$$

Assume the following.

$$\begin{aligned}
& \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\
& \quad (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\
& \quad X1))) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge ((v1_funct_1 X2) \wedge (v1_finseq_1 \\
& \quad X2))) \Rightarrow ((X2 = k7_finseq_1 X0 X1) \Leftrightarrow ((k4_finseq_1 X2 = k2_finseq_1 \\
& \quad (k2_nat_1 (k3_finseq_1 X0) (k3_finseq_1 X1))) \wedge ((\forall X3.(\\
& \quad v7_ordinal1 X3) \Rightarrow ((X3 \in k4_finseq_1 X0) \Rightarrow (k1_funct_1 X2 X3 = k1_funct_1 \\
& \quad X0 X3))) \wedge (\forall X3.(v7_ordinal1 X3) \Rightarrow ((X3 \in k4_finseq_1 X1) \Rightarrow \\
& \quad (k1_funct_1 X2 (k2_nat_1 (k3_finseq_1 X0) X3) = k1_funct_1 X1 X3))))))))) \\
& \hspace{15em} (7)
\end{aligned}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v1_funcop_1 \\
& \quad X0) \Leftrightarrow (\forall X1.(X1 \in k9_xtuple_0 X0) \Rightarrow ((v1_relat_1 (k1_funct_1 \\
& \quad X0 X1)) \wedge (v1_funct_1 (k1_funct_1 X0 X1)))))) \\
& \hspace{15em} (8)
\end{aligned}$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \hspace{15em} (9)$$

Theorem 1

$$\begin{aligned}
& \forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge (v1_finseq_1 X0))) \Rightarrow \\
& \quad (\forall X1.((v1_relat_1 X1) \wedge ((v1_funct_1 X1) \wedge (v1_finseq_1 \\
& \quad X1))) \Rightarrow ((v1_funcop_1 (k7_finseq_1 X0 X1)) \Rightarrow ((v1_funcop_1 X0) \wedge \\
& \quad (v1_funcop_1 X1))))))
\end{aligned}$$