

t23_nat_1 (TM-
PhX8GNzYW98amJ9bRppyV2WjtKtXteBQA)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_2 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\neg (\neg r1_xreal_0 (k1_nat_1 X1 np_1) X0) \wedge ((r1_xreal_0 X1 X0) \wedge (X0 \neq X1)))) \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((\neg r1_xreal_0 np_1 X0) \Rightarrow (X0 = k6_numbers)) \quad (2)$$

Assume the following.

$$((v2_xreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (3)$$

Assume the following.

$$k2_xcmplx_0 np_1 np_1 = np_2 \quad (4)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (m1_subset_1 X1 k5_numbers)) \Rightarrow (k1_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (6)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (7)$$

Theorem 1

$$\forall X0.(v7_ordinal1\ X0) \Rightarrow (\neg(\neg r1_xreal_0\ np_2\ X0) \wedge ((X0 \neq k6_numbers) \wedge (X0 \neq np_1)))$$