

t23_polyeq_4

(TMb29KMH3Cx3YCPrfmUNJ1TtUwKd7hGbKso)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k4_power : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k5_power : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_power : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge (X0 \neq np_1) \wedge (k5_power X0 X0 \neq np_1))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k1_numbers) \wedge (m1_subset_1 X1 k1_numbers)) \Rightarrow (k4_power X0 X1 = k3_power X0 X1) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\neg(\neg r1_xxreal_0 X0 k6_numbers) \wedge ((X0 \neq np_1) \wedge ((\neg r1_xxreal_0 X1 k6_numbers) \wedge (\neg \forall X2.(v1_xreal_0 X2) \Rightarrow ((X2 = k5_power X0 X1) \Leftrightarrow (k3_power X0 X2 = X1)))))))) \quad (3)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (4)$$

Theorem 1

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 k1_numbers) \Rightarrow ((k4_power X0 X1 = X0) \Rightarrow ((r1_xxreal_0 X0 k6_numbers) \vee ((X0 = np_1) \vee (X1 = np_1))))))$$