

t23\_relset\_1 (TM-  
RUKR6jwouuWd4rkULAFVigAnnYWs63pD5)

October 27, 2020

Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k7\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k8\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k8\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k10\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k7\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 \\ & (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((k7\_relset\_1 X0 X1 X2 X0 = k2\_relset\_1 X1 \\ & X2) \wedge (k8\_relset\_1 X0 X1 X2 X1 = k1\_relset\_1 X0 X2)) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. (v1\_relat\_1 X0) \Rightarrow (k8\_relat\_1 X0 (k10\_xtuple\_0 X0) = k9\_xtuple\_0 X0) \quad (2)$$

Assume the following.

$$\forall X0. (v1\_relat\_1 X0) \Rightarrow (k7\_relat\_1 X0 (k9\_xtuple\_0 X0) = k10\_xtuple\_0 X0) \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. (m1\_subset\_1 X2 ( \\ & k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (k8\_relset\_1 X0 X1 X2 X3 = k8\_relat\_1 \\ & X2 X3) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. (m1\_subset\_1 X2 ( \\ & k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (k7\_relset\_1 X0 X1 X2 X3 = k7\_relat\_1 \\ & X2 X3) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge(v5\_relat\_1 X1 X0))\Rightarrow(k2\_relset\_1 X0 X1 = k10\_xtuple\_0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1)\wedge(v4\_relat\_1 X1 X0))\Rightarrow(k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow((v4\_relat\_1 X2 X0)\wedge(v5\_relat\_1 X2 X1)) \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))\Rightarrow(v1\_relat\_1 X2) \quad (9)$$

**Theorem 1**

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X1 X0)))\Rightarrow((k7\_relset\_1 X1 X0 X2 (k8\_relset\_1 X1 X0 X2 X0) = k2\_relset\_1 X0 X2)\wedge(k8\_relset\_1 X1 X0 X2 (k7\_relset\_1 X1 X0 X2 X1) = k1\_relset\_1 X1 X2))$$