

t23_taylor_1 (TMbn-
wYKmb9BA1BcrwZGh8xP3RJxAeBns5jK)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_taylor_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k5_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $r2_fdiff_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_seqfunc : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_taylor_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\ & (r1_xxreal_0 X0 X2)))) \end{aligned} \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow (((r1_xxreal_0 \\ & X0 X1) \wedge (r1_xxreal_0 X2 X3)) \Rightarrow (r1_xxreal_0 (k6_xcmplx_0 X0 X3) (\\ & k6_xcmplx_0 X1 X2)))))) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (3)$$

Assume the following.

$$r1_xxreal_0 np_1 np_1 \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(m1_subset_1 X1 k1_numbers))\Rightarrow (k5_real_1 X0 X1 = k6_xcmplx_0 X0 X1) \quad (6)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0)\wedge(v1_xreal_0 X1))\Rightarrow(v1_xreal_0 (k6_xcmplx_0 X0 X1)) \quad (8)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (9)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (10)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (11)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_funct_1 X0)\wedge(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ k1_numbers k1_numbers))))\Rightarrow(\forall X1.(v7_ordinal1 X1)\Rightarrow(\forall X2. \\ (m1_subset_1 X2 (k1_zfmisc_1 k1_numbers))\Rightarrow((r1_taylor_1 X0 X1 \\ X2)\Leftrightarrow(\forall X3.(m2_subset_1 X3 k1_numbers k5_numbers)\Rightarrow((r1_xxreal_0 \\ X3 (k5_real_1 X1 np_1))\Rightarrow(r2_fdiff_1 (k1_seqfunc k1_numbers k1_numbers \\ (k5_taylor_1 X0 X2) X3) X2)))))) \end{aligned} \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (13)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xxreal_0 X0) \quad (14)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_xxreal_0 X0) \quad (15)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0)\Rightarrow(v1_xreal_0 X0) \quad (16)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_funct_1 X0)\wedge(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ k1_numbers k1_numbers))))\Rightarrow(\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 \\ k1_numbers))\Rightarrow(\forall X2.(v7_ordinal1 X2)\Rightarrow((r1_taylor_1 X0 \\ X2 X1)\Rightarrow(\forall X3.(v7_ordinal1 X3)\Rightarrow((r1_xxreal_0 X3 X2)\Rightarrow(r1_taylor_1 \\ X0 X3 X1)))))) \end{aligned}$$