

t23_tdlat_3
(TMHQ4XNNARAHxjHtRpEGCMfakRCDwasPRH5)

October 27, 2020

Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $v3_tdlat_3 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $v3_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_pre_topc : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v4_pre_topc : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(l1_pre_topc X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (((v4_pre_topc X1 X0) \Rightarrow (k2_pre_topc X0 X1 = X1)) \wedge (((v2_pre_topc X0) \wedge (k2_pre_topc X0 X1 = X1)) \Rightarrow (v4_pre_topc X1 X0)))) \quad (1)$$

Assume the following.

$$\forall X0.((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow ((v3_tdlat_3 X0) \Leftrightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((v3_pre_topc X1 X0) \Rightarrow (v4_pre_topc X1 X0)))) \quad (2)$$

Theorem 1

$$\forall X0.((v2_pre_topc X0) \wedge (l1_pre_topc X0)) \Rightarrow ((v3_tdlat_3 X0) \Leftrightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow ((v3_pre_topc X1 X0) \Rightarrow (k2_pre_topc X0 X1 = X1))))$$