

## t23\_tex\_2

(TMPnX7YZZjrTdt9uG4Fpm1RmNumuYYQEvvQ3)

October 27, 2020

Let  $l1\_pre\_topc : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $u1\_struct\_0 : \iota \Rightarrow \iota$  be given. Let  $v2\_tex\_2 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $r1\_tarski : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k3\_xboole\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\begin{aligned} \forall X0.(l1\_pre\_topc\ X0) \Rightarrow (\forall X1.(m1\_subset\_1\ X1\ (k1\_zfmisc\_1 \\ (u1\_struct\_0\ X0))) \Rightarrow (\forall X2.(m1\_subset\_1\ X2\ (k1\_zfmisc\_1 \\ (u1\_struct\_0\ X0))) \Rightarrow ((r1\_tarski\ X2\ X1) \wedge (v2\_tex\_2\ X1\ X0)) \Rightarrow (v2\_tex\_2 \\ X2\ X0)))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.\forall X1.r1\_tarski\ (k3\_xboole\_0\ X0\ X1)\ X0 \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1\ X2\ (k1\_zfmisc\_1\ X0)) \Rightarrow (k9\_subset\_1\ X0\ X1\ X2 = k3\_xboole\_0\ X1\ X2) \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1\ X2\ (k1\_zfmisc\_1\ X0)) \Rightarrow (m1\_subset\_1\ (k9\_subset\_1\ X0\ X1\ X2)\ (k1\_zfmisc\_1\ X0)) \tag{4}$$

Assume the following.

$$\forall X0.\forall X1.k3\_xboole\_0\ X0\ X1 = k3\_xboole\_0\ X1\ X0 \tag{5}$$

### Theorem 1

$$\begin{aligned} \forall X0.(l1\_pre\_topc\ X0) \Rightarrow (\forall X1.(m1\_subset\_1\ X1\ (k1\_zfmisc\_1 \\ (u1\_struct\_0\ X0))) \Rightarrow (\forall X2.(m1\_subset\_1\ X2\ (k1\_zfmisc\_1 \\ (u1\_struct\_0\ X0))) \Rightarrow ((v2\_tex\_2\ X1\ X0) \vee (v2\_tex\_2\ X2\ X0)) \Rightarrow (v2\_tex\_2 \\ (k9\_subset\_1\ (u1\_struct\_0\ X0)\ X1\ X2)\ X0)))) \end{aligned}$$