

t23_xreal_1

(TMN4Ko145mSahJTAUBvEn22dfcvUEGn8QgY)

October 27, 2020

Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 X0 (k2_xcmplx_0 X1 X2)) \Leftrightarrow (r1_xxreal_0 \\ & (k6_xcmplx_0 X0 X1) X2)))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((v1_xcmplx_0 X0) \wedge ((v1_xcmplx_0 \\ & X1) \wedge (v1_xcmplx_0 X2))) \Rightarrow (k2_xcmplx_0 (k2_xcmplx_0 X0 X1) X2 = k2_xcmplx_0 \\ & X0 (k2_xcmplx_0 X1 X2)) \end{aligned} \tag{2}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xreal_0 X0) \wedge (v1_xreal_0 X1)) \Rightarrow (v1_xreal_0 (k2_xcmplx_0 X0 X1)) \tag{3}$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0) \wedge (v1_xcmplx_0 X1)) \Rightarrow (k2_xcmplx_0 X0 X1 = k2_xcmplx_0 X1 X0) \tag{4}$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (v1_xcmplx_0 X0) \tag{5}$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow ((r1_xxreal_0 (\\ & k6_xcmplx_0 X0 X1) (k2_xcmplx_0 X2 X3)) \Rightarrow (r1_xxreal_0 (k6_xcmplx_0 \\ & X0 X3) (k2_xcmplx_0 X2 X1)))))) \end{aligned}$$