

t243_xxreal_1

(TMW7jHzDErVvj1JfRV1tSSJNP1Jcf4VkoPM)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xxreal_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xxreal_0 : \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow (\forall X3.(v1_xxreal_0 X3) \Rightarrow ((r1_xxreal_0 \\ & X2 X3) \Rightarrow ((r1_xxreal_0 X1 X0) \vee (r1_tarski (k2_xxreal_1 X1 X2) (k3_xxreal_1 \\ & X0 X3))))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (r1_xxreal_0 X0 k1_xxreal_0) \tag{2}$$

Assume the following.

$$v1_xxreal_0 k1_xxreal_0 \tag{3}$$

Theorem 1

$$\begin{aligned} & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xxreal_0 X2) \Rightarrow ((\neg r1_xxreal_0 X1 X0) \Rightarrow (r1_tarski (k2_xxreal_1 \\ & X1 X2) (k3_xxreal_1 X0 k1_xxreal_0)))))) \end{aligned}$$