

t24_catalan2
(TMX8m122CsEqn7oBcmaDWyhZtVHSPMNBByfW)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k5_card_1 : \iota \Rightarrow \iota$ be given. Let $k2_catalan2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $k1_card_1 : \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_numbers : \iota$ be given. Let $k8_funcop_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. k1_card_1 (k1_tarski X0) = np_1 \quad (1)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (k2_catalan2 X0 k6_numbers = k6_domain_1 (k1_zfmisc_1 (k2_zfmisc_1 X0 k5_numbers)) (k8_funcop_1 k5_numbers X0 k6_numbers)) \quad (2)$$

Assume the following.

$$m1_subset_1 k1_xboole_0 k4_ordinal1 \quad (3)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge (m1_subset_1 X1 X0)) \Rightarrow (k6_domain_1 X0 X1 = k1_tarski X1) \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$\forall X0.(v1_finset_1 X0) \Rightarrow (k5_card_1 X0 = k1_card_1 X0) \quad (7)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (8)$$

Assume the following.

$$\forall X0. \neg v1_xboole_0 (k1_zfmisc_1 X0) \quad (9)$$

Assume the following.

$$\forall X0.v1_finset_1 (k1_tarski X0) \quad (10)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.((\neg v1_xboole_0 X0) \wedge (m1_subset_1 \\ & X2 X0)) \Rightarrow ((v1_funct_1 (k8_funcop_1 X0 X1 X2)) \wedge ((v1_funct_2 (k8_funcop_1 \\ & X0 X1 X2) X1 X0) \wedge (m1_subset_1 (k8_funcop_1 X0 X1 X2) (k1_zfmisc_1 \\ & (k2_zfmisc_1 X1 X0)))))) \end{aligned} \quad (11)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (k5_card_1 (k2_catalan2 X0 k6_numbers) = np_1)$$