

t24_comseq_2 (TMXuK-
WHosKQuq8BqiSZn73eV6rx5WJGkcY8)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k2_numbers : \iota$ be given. Let $v2_comseq_2 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_comseq_2 : \iota \Rightarrow \iota$ be given. Let $k2_comseq_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k31_valued_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_complex1 : \iota \Rightarrow \iota$ be given. Let $k15_complex1 : \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $v1_membered : \iota \Rightarrow o$ be given. Let $k30_valued_1 : \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k15_complex1 (k4_xcmplx_0 X0) = k10_complex1 (k15_complex1 X0)) \quad (1)$$

Assume the following.

$$\forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k2_numbers) \wedge ((v2_comseq_2 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k2_numbers)))))) \Rightarrow (k3_comseq_2 (k31_valued_1 k5_numbers k2_numbers X0) = k10_complex1 (k3_comseq_2 X0)) \quad (2)$$

Assume the following.

$$\forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k2_numbers) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k2_numbers)))))) \Rightarrow ((v2_comseq_2 X0) \Rightarrow (k3_comseq_2 (k2_comseq_2 k5_numbers X0) = k15_complex1 (k3_comseq_2 X0))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.((v1_membered X1) \wedge ((v1_funct_1 X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow (k31_valued_1 X0 X1 X2 = k30_valued_1 X2) \quad (4)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k2_numbers) \Rightarrow (k10_complex1 X0 = k4_xcmplx_0 X0) \quad (5)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k2_numbers) \wedge \\ ((v2_comseq_2 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ k5_numbers k2_numbers)))))) \Rightarrow ((v1_funct_1 (k30_valued_1 X0)) \wedge \\ ((v1_funct_2 (k30_valued_1 X0) k5_numbers k2_numbers) \wedge (v2_comseq_2 \\ (k30_valued_1 X0)))) \end{aligned} \quad (6)$$

Assume the following.

$$v1_membered k2_numbers \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k2_numbers) \wedge \\ (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k2_numbers)))))) \Rightarrow \\ (m1_subset_1 (k3_comseq_2 X0) k2_numbers) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.((v1_membered X1) \wedge ((v1_funct_1 \\ X2) \wedge (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))))) \Rightarrow ((v1_funct_1 \\ (k31_valued_1 X0 X1 X2)) \wedge (m1_subset_1 (k31_valued_1 X0 X1 X2) (\\ k1_zfmisc_1 (k2_zfmisc_1 X0 k2_numbers)))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k2_numbers) \Rightarrow (v1_xcmplx_0 X0) \quad (10)$$

Theorem 1

$$\begin{aligned} \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k2_numbers) \wedge \\ ((v2_comseq_2 X0) \wedge (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 \\ k5_numbers k2_numbers)))))) \Rightarrow (k3_comseq_2 (k2_comseq_2 k5_numbers \\ (k31_valued_1 k5_numbers k2_numbers X0)) = k10_complex1 (k15_complex1 \\ (k3_comseq_2 X0))) \end{aligned}$$