

t24_comseq_3
(TMZXr8fmb6fW8hzxwnVXzR88NL4CcKxde7r)

October 27, 2020

Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k2_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k8_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_complex1 : \iota$ be given. Let $k10_comseq_3 : \iota \Rightarrow \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $k2_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_series_1 : \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_valued_0 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_partfun1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k2_xcmplx_0 X0 \ k6_numbers = X0) \quad (1)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 \ np_1) \wedge (m2_subset_1 \ np_1 \ k1_numbers \ k5_numbers)) \wedge \\ & ((m1_subset_1 \ np_1 \ k5_numbers) \wedge (m1_subset_1 \ np_1 \ k1_numbers)) \end{aligned} \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge \\ & (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2. (m2_subset_1 \\ & X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \end{aligned} \quad (3)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (((v1_funct_1 X1) \wedge ((v1_funct_2 \\ & X1 \ k5_numbers X0) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \ k5_numbers \\ & X0)))))) \wedge (v7_ordinal1 X2)) \Rightarrow (k8_nat_1 X0 X1 X2 = k1_funct_1 X1 X2) \end{aligned} \quad (4)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$k5_complex1 = k1_xboole_0 \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(k2_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (8)$$

Assume the following.

$$\forall X0.((v1_funct_1 X0)\wedge((v1_funct_2 X0 k5_numbers k2_numbers)\wedge(m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k2_numbers))))\Rightarrow(k10_comseq_3 X0 = k2_series_1 X0) \quad (9)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X0)\wedge((v1_funct_1 X0)\wedge(v1_valued_0 X0)))\Rightarrow(v1_xcmplx_0 (k1_funct_1 X0 X1)) \quad (11)$$

Assume the following.

$$\neg v1_xboole_0 k2_numbers \quad (12)$$

Assume the following.

$$\neg v1_xboole_0 k1_numbers \quad (13)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(v7_ordinal1 X1))\Rightarrow(v7_ordinal1 (k2_xcmplx_0 X0 X1)) \quad (14)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k5_numbers)\wedge(v7_ordinal1 X1))\Rightarrow(m2_subset_1 (k2_nat_1 X0 X1) k1_numbers k5_numbers) \quad (16)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k2_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k2_numbers)))))) \Rightarrow \\ & ((v1_funct_1 (k10_comseq_3 X0)) \wedge ((v1_funct_2 (k10_comseq_3 \\ & X0) k5_numbers k2_numbers) \wedge (m1_subset_1 (k10_comseq_3 X0) (k1_zfmisc_1 \\ & (k2_zfmisc_1 k5_numbers k2_numbers)))))) \end{aligned} \quad (17)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_relat_1 X0) \wedge ((v4_relat_1 X0 k5_numbers) \wedge ((v1_funct_1 \\ & X0) \wedge ((v1_partfun1 X0 k5_numbers) \wedge (v1_valued_0 X0)))))) \Rightarrow (\forall X1. \\ & ((v1_relat_1 X1) \wedge ((v4_relat_1 X1 k5_numbers) \wedge ((v1_funct_1 X1) \wedge \\ & ((v1_partfun1 X1 k5_numbers) \wedge (v1_valued_0 X1)))))) \Rightarrow ((X1 = k2_series_1 \\ & X0) \Leftrightarrow ((k1_funct_1 X1 k6_numbers = k1_funct_1 X0 k6_numbers) \wedge (\forall X2. \\ & (m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow (k1_funct_1 X1 (k2_nat_1 \\ & X2 np_1) = k2_xcmplx_0 (k1_funct_1 X1 X2) (k1_funct_1 X0 (k2_nat_1 \\ & X2 np_1)))))) \end{aligned} \quad (18)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (19)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. (\neg v1_xboole_0 X1) \Rightarrow (\forall X2. (m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow ((v1_funct_2 X2 X0 X1) \Rightarrow (\\ & v1_partfun1 X2 X0))) \end{aligned} \quad (20)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k2_zfmisc_1 X0 X1))) \Rightarrow ((v4_relat_1 X2 X0) \wedge (v5_relat_1 X2 X1)) \end{aligned} \quad (21)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v5_relat_1 X0 k2_numbers)) \Rightarrow ((v1_relat_1 X0) \wedge (v1_valued_0 X0)) \quad (22)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (23)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((v1_funct_1 X0) \wedge ((v1_funct_2 X0 k5_numbers k2_numbers) \wedge \\ & (m1_subset_1 X0 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers k2_numbers)))))) \Rightarrow \\ & ((\forall X1. (m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (k8_nat_1 \\ & k2_numbers X0 X1 = k5_complex1)) \Rightarrow (\forall X1. (m2_subset_1 X1 k1_numbers \\ & k5_numbers) \Rightarrow (k8_nat_1 k2_numbers (k10_comseq_3 X0) X1 = k5_complex1))) \end{aligned}$$