

t24_euclid_3

(TMSMGHeA7pHUZi7T6aFUcb8vHBjxvnwFttd)

October 27, 2020

Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k12_euclid : \iota \Rightarrow \iota$ be given. Let $k1_euclid_3 : \iota \Rightarrow \iota$ be given. Let $k7_square_1 : \iota \Rightarrow \iota$ be given. Let $k7_real_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_square_1 : \iota \Rightarrow \iota$ be given. Let $k3_complex1 : \iota \Rightarrow \iota$ be given. Let $k4_complex1 : \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k19_euclid : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k2_complex1 : \iota \Rightarrow \iota$ be given. Let $k1_complex1 : \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow ((k17_euclid (k19_euclid X0 X1) = X0) \wedge (k18_euclid (k19_euclid X0 X1) = X1))) \quad (1)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\ & ((k12_euclid X0 = k7_square_1 (k7_real_1 (k5_square_1 (k17_euclid X0)) (k5_square_1 (k18_euclid X0)))) \wedge (k5_square_1 (k12_euclid X0) = k7_real_1 (k5_square_1 (k17_euclid X0)) (k5_square_1 (k18_euclid X0)))) \end{aligned} \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k4_complex1 X0 = k2_complex1 X0) \quad (3)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k3_complex1 X0 = k1_complex1 X0) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (m1_subset_1 (k4_complex1 X0) k1_numbers) \quad (5)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (m1_subset_1 (k3_complex1 X0) k1_numbers) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (m1_subset_1 (k1_euclid_3 X0) (u1_struct_0 (k15_euclid np_2))) \quad (7)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k1_euclid_3 X0 = k19_euclid (k3_complex1 X0) (k4_complex1 X0)) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers) \Rightarrow (v1_xreal_0 X0) \quad (9)$$

Theorem 1

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k12_euclid (k1_euclid_3 X0) = k7_square_1 (k7_real_1 (k5_square_1 (k3_complex1 X0)) (k5_square_1 (k4_complex1 X0))))$$