

t24_jgraph_7

(TMc43xBwnmLskPg7duRFJopygW8pTFNCxiN)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $k17_euclid : \iota \Rightarrow \iota$ be given. Let $k18_euclid : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $r1_jordan17 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_sppol_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_jordan6 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Assume the following.

$$\begin{aligned}
 & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\
 & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\
 & (\forall X2.(v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow (\forall X4. \\
 & (v1_xreal_0 X4) \Rightarrow (\forall X5.(v1_xreal_0 X5) \Rightarrow (((k18_euclid X0 = \\
 & X5) \wedge ((k18_euclid X1 = X5) \wedge ((r1_xxreal_0 X2 (k17_euclid X0)) \wedge (\\
 & r1_xxreal_0 (k17_euclid X1) X3)))) \Rightarrow ((r1_xxreal_0 X3 X2) \vee ((r1_xxreal_0 \\
 & X5 X4) \vee ((r1_xxreal_0 (k17_euclid X1) (k17_euclid X0)) \vee (r1_jordan6 \\
 & (k1_sppol_2 X2 X3 X4 X5) X0 X1))))))))))
 \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\
 & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2))) \Rightarrow \\
 & (\forall X2.(v1_xreal_0 X2) \Rightarrow (\forall X3.(v1_xreal_0 X3) \Rightarrow (((\\
 & r1_xxreal_0 X3 (k18_euclid X0)) \wedge ((r1_xxreal_0 (k18_euclid X0) \\
 & (k18_euclid X1)) \wedge ((r1_xxreal_0 (k17_euclid X0) (k17_euclid X1)) \wedge \\
 & (r1_xxreal_0 (k17_euclid X1) X2)))) \Rightarrow ((r1_xxreal_0 X2 (k17_euclid \\
 & X0)) \vee ((r1_xxreal_0 (k18_euclid X1) X3) \vee (r1_jordan6 (k1_sppol_2 \\
 & (k17_euclid X0) X2 X3 (k18_euclid X1)) X0 X1))))))
 \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
 & \forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2. \\
 & (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow \\
 & (r1_xxreal_0 X0 X2))))
 \end{aligned} \tag{3}$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.\forall X3.((v1_xreal_0 X0)\wedge \\ & ((v1_xreal_0 X1)\wedge((v1_xreal_0 X2)\wedge(v1_xreal_0 X3))))\Rightarrow(m1_subset_1 \\ & (k1_sppol_2 X0 X1 X2 X3) (k1_zfmisc_1 (u1_struct_0 (k15_euclid \\ & np_2)))) \end{aligned} \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2)))\Rightarrow \\ & (m1_subset_1 (k17_euclid X0) k1_numbers) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (k1_zfmisc_1 (u1_struct_0 (k15_euclid \\ & np_2))))\Rightarrow(\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid \\ & np_2)))\Rightarrow(\forall X2.(m1_subset_1 X2 (u1_struct_0 (k15_euclid \\ & np_2)))\Rightarrow(\forall X3.(m1_subset_1 X3 (u1_struct_0 (k15_euclid \\ & np_2)))\Rightarrow(\forall X4.(m1_subset_1 X4 (u1_struct_0 (k15_euclid \\ & np_2))))\Rightarrow((r1_jordan17 X0 X1 X2 X3 X4)\Leftrightarrow(\neg(\neg(r1_jordan6 X0 X1 X2)\wedge \\ & ((r1_jordan6 X0 X2 X3)\wedge(r1_jordan6 X0 X3 X4))\wedge(\neg(r1_jordan6 \\ & X0 X2 X3)\wedge(r1_jordan6 X0 X3 X4)\wedge(r1_jordan6 X0 X4 X1)))\wedge(\neg(r1_jordan6 \\ & X0 X3 X4)\wedge(r1_jordan6 X0 X4 X1)\wedge(r1_jordan6 X0 X1 X2)))\wedge(\neg(r1_jordan6 \\ & X0 X4 X1)\wedge(r1_jordan6 X0 X1 X2)\wedge(r1_jordan6 X0 X2 X3)))))))))) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v1_xxreal_0 X0)\wedge(v1_xxreal_0 X1))\Rightarrow(\\ & (r1_xxreal_0 X0 X1)\vee(r1_xxreal_0 X1 X0)) \end{aligned} \quad (7)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xxreal_0 X0) \quad (8)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (9)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m1_subset_1 X0 (u1_struct_0 (k15_euclid np_2)))\Rightarrow \\ & (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid np_2)))\Rightarrow \\ & (\forall X2.(m1_subset_1 X2 (u1_struct_0 (k15_euclid np_2)))\Rightarrow \\ & (\forall X3.(m1_subset_1 X3 (u1_struct_0 (k15_euclid np_2)))\Rightarrow \\ & (\forall X4.(v1_xreal_0 X4)\Rightarrow(\forall X5.(v1_xreal_0 X5)\Rightarrow(\forall X6. \\ & (v1_xreal_0 X6)\Rightarrow(\forall X7.(v1_xreal_0 X7)\Rightarrow(((k17_euclid X0 = \\ & X4)\wedge((k18_euclid X1 = X7)\wedge((k18_euclid X2 = X7)\wedge((k18_euclid X3 = \\ & X7)\wedge((r1_xxreal_0 X6 (k18_euclid X0))\wedge((r1_xxreal_0 (k18_euclid \\ & X0) X7)\wedge((r1_xxreal_0 X4 (k17_euclid X1))\wedge(r1_xxreal_0 (k17_euclid \\ & X3) X5))))))))))\Rightarrow((r1_xxreal_0 X5 X4)\vee((r1_xxreal_0 X7 X6)\vee((r1_xxreal_0 \\ & (k17_euclid X2) (k17_euclid X1))\vee((r1_xxreal_0 (k17_euclid X3) \\ & (k17_euclid X2))\vee(r1_jordan17 (k1_sppol_2 X4 X5 X6 X7) X0 X1 X2 X3)))))))))) \end{aligned}$$