

t24_orders_2 (TMEnDqxs- FvpotcMKY6aSNUxi5w8ehHDbpKq)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v3_orders_2 : \iota \Rightarrow o$ be given. Let $v4_orders_2 : \iota \Rightarrow o$ be given. Let $v5_orders_2 : \iota \Rightarrow o$ be given. Let $l1_orders_2 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k3_orders_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r2_orders_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_orders_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_domain_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.((\neg v2_struct_0 X0) \wedge ((v3_orders_2 X0) \wedge ((v4_orders_2 \\ X0) \wedge ((v5_orders_2 X0) \wedge (l1_orders_2 X0)))))) \Rightarrow (\forall X1.(m1_subset_1 \\ X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 \\ X0)) \Rightarrow ((r2_orders_2 X0 X1 X2) \Leftrightarrow (X1 \in k2_orders_2 X0 (k6_domain_1 \\ (u1_struct_0 X0) X2)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 X0)) \Rightarrow (k9_subset_1 X0 X1 X2 = k3_xboole_0 X1 X2) \tag{2}$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (X2 = k3_xboole_0 X0 X1) \Leftrightarrow (\forall X3. (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (X3 \in X1))) \tag{3}$$

Assume the following.

$$\begin{aligned} \forall X0.((\neg v2_struct_0 X0) \wedge ((v3_orders_2 X0) \wedge ((v4_orders_2 \\ X0) \wedge ((v5_orders_2 X0) \wedge (l1_orders_2 X0)))))) \Rightarrow (\forall X1.(m1_subset_1 \\ X1 (k1_zfmisc_1 (u1_struct_0 X0))) \Rightarrow (\forall X2.(m1_subset_1 \\ X2 (u1_struct_0 X0)) \Rightarrow (k3_orders_2 X0 X1 X2 = k9_subset_1 (u1_struct_0 \\ X0) (k2_orders_2 X0 (k6_domain_1 (u1_struct_0 X0) X2)) X1))) \end{aligned} \tag{4}$$

Theorem 1

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v3_orders_2 X0) \wedge ((v4_orders_2 \\ & X0) \wedge ((v5_orders_2 X0) \wedge (l1_orders_2 X0)))))) \Rightarrow (\forall X1.(m1_subset_1 \\ & X1 (u1_struct_0 X0)) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 \\ & X0)) \Rightarrow (\forall X3.(m1_subset_1 X3 (k1_zfmisc_1 (u1_struct_0 X0)) \Rightarrow \\ & ((X1 \in k3_orders_2 X0 X3 X2) \Leftrightarrow ((r2_orders_2 X0 X1 X2) \wedge (X1 \in X3))))))) \end{aligned}$$