

# t24\_partfun2

## (TMZ66hhUBjbyGz6hvGdSFe7aeKwNptPzss2)

October 27, 2020

Let  $v1\_xboole\_0 : \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k9\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $k7\_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((X0 \in k9\_xtuple\_0 X1) \Rightarrow (k9\_relat\_1 X1 X0 = k1\_tarski (k1\_funct\_1 X1 X0))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge ((v5\_relat\_1 X1 X0) \wedge (v1\_funct\_1 X1))) \Rightarrow (\forall X2. (X2 \in k9\_xtuple\_0 X1) \Rightarrow (k7\_partfun1 X0 X1 X2 = k1\_funct\_1 X1 X2)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v4\_relat\_1 X2 X0) \wedge (v5\_relat\_1 X2 X1)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_relat\_1 X2) \quad (5)$$

### Theorem 1

$$\forall X0. (\neg v1\_xboole\_0 X0) \Rightarrow (\forall X1. (\neg v1\_xboole\_0 X1) \Rightarrow (\forall X2. (m1\_subset\_1 X2 X0) \Rightarrow (\forall X3. ((v1\_funct\_1 X3) \wedge (m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1)))) \Rightarrow ((X2 \in k1\_relset\_1 X0 X3) \Rightarrow (k9\_relat\_1 X3 X2 = k1\_tarski (k7\_partfun1 X1 X3 X2)))))))$$