

t24\_pepin (TMQXVas-  
RAZp6gMpJQmHUyC3W8by9UTsp77C)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_newton : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $np\_2 : \iota$  be given. Let  $k3\_square\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_xcmplx\_0 : \iota \Rightarrow o$  be given. Let  $k3\_xcmplx\_0 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xcmplx\_0 X0) \Rightarrow ((k1\_newton X0 np\_2 = k3\_xcmplx\_0 X0 X0) \wedge (k3\_square\_1 X0 = k1\_newton X0 np\_2)) \quad (1)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (v1\_xcmplx\_0 X0) \quad (2)$$

**Theorem 1**  $\forall X0.(v7\_ordinal1 X0) \Rightarrow (k1\_newton X0 np\_2 = k3\_square\_1 X0)$ .