

t24_prob_3 (TMHJwkKiKuXG- GvZqtCptEmh63xyoYbo9pwn)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_prob_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $k9_setfam_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_prob_3 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.\forall X2.((v1_funct_1 \\ & X2) \wedge ((v1_funct_2 X2 X1) \wedge (k5_numbers (k9_setfam_1 X1)) \wedge (m1_subset_1 \\ & X2 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k9_setfam_1 X1)))))) \Rightarrow \\ & (r1_tarski (k1_funct_1 X2 X0) (k1_funct_1 (k2_prob_3 X1 X2) X0))) \end{aligned} \quad (1)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.\forall X2.((\neg v1_xboole_0 \\ & X2) \wedge ((v1_prob_1 X2 X1) \wedge ((v4_prob_1 X2 X1) \wedge (m1_subset_1 X2 (k1_zfmisc_1 \\ & (k1_zfmisc_1 X1)))))) \Rightarrow (\forall X3.((v5_relat_1 X3 X2) \wedge ((v1_funct_1 \\ & X3) \wedge ((v1_funct_2 X3 k5_numbers (k9_setfam_1 X1)) \wedge (m1_subset_1 \\ & X3 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k9_setfam_1 X1)))))) \Rightarrow \\ & (r1_tarski (k1_funct_1 X3 X0) (k1_funct_1 (k2_prob_3 X1 X3) X0)))) \end{aligned}$$