

t24_xxreal_0

(TMTJBYNsVQD3RaR8YR8JeDMYn9t1GjjjxS5)

October 27, 2020

Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_xxreal_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (r1_xxreal_0 (k3_xxreal_0 X0 X1) X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \Rightarrow (k3_xxreal_0 X0 X1 = X0)) \wedge ((\neg r1_xxreal_0 X0 X1) \Rightarrow (k3_xxreal_0 X0 X1 = X1)))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (r1_xxreal_0 X0 X1) \vee (r1_xxreal_0 X1 X0) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xxreal_0 X0) \wedge (v1_xxreal_0 X1)) \Rightarrow (k3_xxreal_0 X0 X1 = k3_xxreal_0 X1 X0) \quad (4)$$

Theorem 1

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow (\forall X2.(v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge ((r1_xxreal_0 X0 X2) \wedge (\forall X3.(v1_xxreal_0 X3) \Rightarrow (((r1_xxreal_0 X3 X1) \wedge (r1_xxreal_0 X3 X2)) \Rightarrow (r1_xxreal_0 X3 X0)))))) \Rightarrow (X0 = k3_xxreal_0 X1 X2))))$$