

t25_analort
(TMUfUcmumkL69TvrvtvKjkQgAYnJiun98He7)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_analort : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_analort : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_analof : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} & \forall X0.((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 \\ & X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge (l1_rlvect_1 \\ & X0)))))))))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow \\ & (\forall X2.(m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 \\ & X3 (u1_struct_0 X0)) \Rightarrow (\forall X4.(m1_subset_1 X4 (u1_struct_0 \\ & X0)) \Rightarrow ((k5_algstr_0 X0 X1 X2 = k5_algstr_0 X0 X3 X4) \Rightarrow (r1_analof \\ & X0 X2 X1 X4 X3)))))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. \forall X2. \forall X3. (((\neg v2_struct_0 \\ & X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 X0) \wedge ((v3_rlvect_1 X0) \wedge \\ & ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 \\ & X0) \wedge ((v8_rlvect_1 X0) \wedge (l1_rlvect_1 X0)))))))))) \wedge ((m1_subset_1 \\ & X1 (u1_struct_0 X0)) \wedge ((m1_subset_1 X2 (u1_struct_0 X0)) \wedge (m1_subset_1 \\ & X3 (u1_struct_0 X0)))) \Rightarrow (m1_subset_1 (k2_analort X0 X1 X2 X3) (\\ & u1_struct_0 X0)) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 \\
& X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\
& ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge (l1_rlvect_1 \\
& X0)))))))))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow \\
& (\forall X2.(m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 \\
& X3 (u1_struct_0 X0)) \Rightarrow (\forall X4.(m1_subset_1 X4 (u1_struct_0 \\
& X0)) \Rightarrow (\forall X5.(m1_subset_1 X5 (u1_struct_0 X0)) \Rightarrow (\forall X6. \\
& (m1_subset_1 X6 (u1_struct_0 X0)) \Rightarrow ((r1_analort X0 X1 X2 X3 X4 X5 \\
& X6) \Leftrightarrow (r1_analof X0 (k2_analort X0 X1 X2 X3) (k2_analort X0 X1 X2 X4) \\
& X5 X6))))))))))
\end{aligned} \tag{3}$$

Theorem 1

$$\begin{aligned}
& \forall X0.((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 \\
& X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\
& ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge (l1_rlvect_1 \\
& X0)))))))))) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow \\
& (\forall X2.(m1_subset_1 X2 (u1_struct_0 X0)) \Rightarrow (\forall X3.(m1_subset_1 \\
& X3 (u1_struct_0 X0)) \Rightarrow (\forall X4.(m1_subset_1 X4 (u1_struct_0 \\
& X0)) \Rightarrow (r1_analort X0 X3 X4 X1 X2 (k2_analort X0 X3 X4 X1) (k2_analort \\
& X0 X3 X4 X2))))))
\end{aligned}$$