

t25_binari_4

(TMH8MhFCTCFJWxm71KTxfiW31WsQxKnuzBs)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_series_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $k1_binari_4 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xxreal_0 X0) \Rightarrow (\forall X1.(v1_xxreal_0 X1) \Rightarrow ((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X0)) \Rightarrow (X0 = X1)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (v7_ordinal1 X1)) \Rightarrow (v7_ordinal1 (k1_binari_4 X0 X1)) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\forall X2. \\ & (v7_ordinal1 X2) \Rightarrow ((X2 = k1_binari_4 X0 X1) \Leftrightarrow ((r1_xxreal_0 X1 (k5_series_1 \\ & np_2 X2)) \wedge ((r1_xxreal_0 X0 X2) \wedge (\forall X3.(v7_ordinal1 X3) \Rightarrow \\ & ((r1_xxreal_0 X1 (k5_series_1 np_2 X3)) \wedge (r1_xxreal_0 X0 X3)) \Rightarrow \\ & (r1_xxreal_0 X2 X3)))))))) \quad (3) \end{aligned}$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xxreal_0 X0) \quad (4)$$

Theorem 1

$$\begin{aligned} & \forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\neg \\ & (\neg r1_xxreal_0 X0 (k5_series_1 np_2 X1)) \wedge (r1_xxreal_0 (k1_binari_4 \\ & X1 X0) X1))) \end{aligned}$$