

# t25\_eqrel\_1

(TMG9cPGxNNiQBqJRP9AJkXoeaBL4riof3ZG)

October 27, 2020

Let  $k6\_eqrel\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k6\_partfun1 : \iota \Rightarrow \iota$  be given. Let  $k1\_tarski : \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k9\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k4\_relat\_1 : \iota \Rightarrow \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $v1\_partfun1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((X0 \in k9\_xtuple\_0 X1) \Rightarrow (k9\_relat\_1 X1 X0 = k1\_tarski (k1\_funct\_1 X1 X0))) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow ((X1 = k4\_relat\_1 X0) \Leftrightarrow ((k9\_xtuple\_0 X1 = X0) \wedge (\forall X2. (X2 \in X0) \Rightarrow (k1\_funct\_1 X1 X2 = X2)))) \quad (2)$$

Assume the following.

$$\forall X0. k6\_partfun1 X0 = k4\_relat\_1 X0 \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. \forall X3. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (k6\_eqrel\_1 X0 X1 X2 X3 = k9\_relat\_1 X2 X3) \quad (4)$$

Assume the following.

$$\forall X0. k9\_xtuple\_0 (k4\_relat\_1 X0) = X0 \quad (5)$$

Assume the following.

$$\forall X0. (v1\_relat\_1 (k4\_relat\_1 X0)) \wedge ((v4\_relat\_1 (k4\_relat\_1 X0) X0) \wedge ((v1\_funct\_1 (k4\_relat\_1 X0)) \wedge (v1\_partfun1 (k4\_relat\_1 X0) X0))) \quad (6)$$

Assume the following.

$$\forall X0.(v1\_partfun1 (k6\_partfun1 X0) X0) \wedge (m1\_subset\_1 (k6\_partfun1 X0) (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X0))) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_relat\_1 X2) \quad (8)$$

**Theorem 1**

$$\forall X0.\forall X1.(X1 \in X0) \Rightarrow (k6\_eqrel\_1 X0 X0 (k6\_partfun1 X0) X1 = k1\_tarski X1)$$