

## t25\_finseq\_3

(TMQK8ANQtuhvwBF2E2sNCZiXJKzN27S2hsw)

October 27, 2020

Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_finseq\_1 : \iota \Rightarrow o$  be given. Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_1 : \iota$  be given. Let  $k3\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_finseq\_1 : \iota \Rightarrow \iota$  be given. Let  $k1\_card\_1 : \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $v1\_finset\_1 : \iota \Rightarrow o$  be given. Let  $v1\_card\_1 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $v3\_ordinal1 : \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((X0 \in k2\_finseq\_1 X1) \Leftrightarrow ((r1\_xxreal\_0 np\_1 X0) \wedge (r1\_xxreal\_0 X0 X1)))) \quad (1)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (k3\_finseq\_1 X0 = k1\_card\_1 X0) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1) \quad (3)$$

Assume the following.

$$\forall X0.(v1\_finset\_1 X0) \Rightarrow ((v1\_finset\_1 (k1\_card\_1 X0)) \wedge (v1\_card\_1 (k1\_card\_1 X0))) \quad (4)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (m2\_subset\_1 (k3\_finseq\_1 X0) k1\_numbers k5\_numbers) \quad (5)$$

Assume the following.

$$\forall X0.((v1\_relat\_1 X0) \wedge ((v1\_funct\_1 X0) \wedge (v1\_finseq\_1 X0))) \Rightarrow (\forall X1.(m2\_subset\_1 X1 k1\_numbers k5\_numbers) \Rightarrow ((X1 = k3\_finseq\_1 X0) \Leftrightarrow (k2\_finseq\_1 X1 = k9\_xtuple\_0 X0))) \quad (6)$$

Assume the following.

$$\forall X0.((v3\_ordinal1 X0)\wedge(v1\_finset\_1 X0))\Rightarrow(v7\_ordinal1 X0) \quad (7)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow \\ ((v1\_relat\_1 X0)\wedge((v4\_relat\_1 X0 k5\_numbers)\wedge((v1\_funct\_1 X0)\wedge \\ (v1\_finseq\_1 X0)))) \end{aligned} \quad (8)$$

Assume the following.

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow \\ ((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finset\_1 X0))) \end{aligned} \quad (9)$$

Assume the following.

$$\forall X0.(v1\_card\_1 X0)\Rightarrow(v3\_ordinal1 X0) \quad (10)$$

**Theorem 1**

$$\begin{aligned} \forall X0.((v1\_relat\_1 X0)\wedge((v1\_funct\_1 X0)\wedge(v1\_finseq\_1 X0)))\Rightarrow \\ (\forall X1.(v7\_ordinal1 X1)\Rightarrow((X1 \in k1\_relset\_1 k5\_numbers X0)\Leftrightarrow \\ ((r1\_xxreal\_0 np\_1 X1)\wedge(r1\_xxreal\_0 X1 (k3\_finseq\_1 X0)))) \end{aligned}$$