

t25_funct_1
(TMZdJ5gMJ1fQxRSNHyDoqpfu81ZGn5KdwGD)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v2_funct_1 : \iota \Rightarrow o$ be given. Let $k3_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.(v1_relat_1 X1) \Rightarrow ((r1_tarski (k10_xtuple_0 X0) (k9_xtuple_0 X1)) \Rightarrow (k9_xtuple_0 (k3_relat_1 X0 X1) = k9_xtuple_0 X0))) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (\forall X2.((v1_relat_1 X2) \wedge (v1_funct_1 X2)) \Rightarrow ((X0 \in k9_xtuple_0 X1) \Rightarrow (k1_funct_1 (k3_relat_1 X1 X2) X0 = k1_funct_1 X2 (k1_funct_1 X1 X0)))) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \wedge ((v1_relat_1 X1) \wedge (v1_funct_1 X1))) \Rightarrow ((v1_relat_1 (k3_relat_1 X0 X1) X1) \wedge (v1_funct_1 (k3_relat_1 X0 X1))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.v1_relat_1 (k3_relat_1 X0 X1) \quad (4)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v2_funct_1 X0) \Leftrightarrow (\forall X1.\forall X2.((X1 \in k9_xtuple_0 X0) \wedge ((X2 \in k9_xtuple_0 X0) \wedge (k1_funct_1 X0 X1 = k1_funct_1 X0 X2)))) \Rightarrow (X1 = X2))) \quad (5)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow (\forall X1.((v1_relat_1 X1) \wedge (v1_funct_1 X1)) \Rightarrow (((v2_funct_1 (k3_relat_1 X1 X0)) \wedge (r1_tarski (k10_xtuple_0 X1) (k9_xtuple_0 X0))) \Rightarrow (v2_funct_1 X1)))$$