

t25_glib_000

(TMUhE84PS2mThJZXtw67xzXnkhEtFbrQxCQ)

October 27, 2020

Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finset_1 : \iota \Rightarrow o$ be given. Let $v1_glib_000 : \iota \Rightarrow o$ be given. Let $v2_glib_000 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k15_glib_000 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\neg(k6_numbers \neq X0) \wedge (r1_xxreal_0 X0 k6_numbers)) \quad (2)$$

Assume the following.

$$\forall X0.(v1_xcmplx_0 X0) \Rightarrow (k2_xcmplx_0 X0 k6_numbers = X0) \quad (3)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow ((\neg r1_xxreal_0 (k1_nat_1 X1 np_1) X0) \Leftrightarrow (r1_xxreal_0 X0 X1))) \quad (4)$$

Assume the following.

$$((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \quad (5)$$

Assume the following.

$$(m2_subset_1 np_0 k1_numbers k5_numbers) \wedge ((m1_subset_1 np_0 k5_numbers) \wedge (m1_subset_1 np_0 k1_numbers)) \quad (6)$$

Assume the following.

$$v1_xboole_0 \text{ np_}0 \quad (7)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (8)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0)\wedge(m1_subset_1 X1 k5_numbers))\Rightarrow (k1_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \quad (10)$$

Assume the following.

$$\forall X0.((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge((v1_finset_1 X0)\wedge((v1_glib_000 X0)\wedge(v2_glib_000 X0))))))\Rightarrow ((\neg v1_xboole_0 (k15_glib_000 X0))\wedge(m1_subset_1 (k15_glib_000 X0) k5_numbers)) \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xcmplx_0 X0)\wedge(v1_xcmplx_0 X1))\Rightarrow (k2_xcmplx_0 X0 X1 = k2_xcmplx_0 X1 X0) \quad (12)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1)\Rightarrow(v7_ordinal1 X0) \quad (13)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0)\Rightarrow(v1_xcmplx_0 X0) \quad (14)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k1_numbers)\Rightarrow(v1_xreal_0 X0) \quad (15)$$

Theorem 1

$$\forall X0.((v1_relat_1 X0)\wedge((v4_relat_1 X0 k5_numbers)\wedge((v1_funct_1 X0)\wedge((v1_finset_1 X0)\wedge((v1_glib_000 X0)\wedge(v2_glib_000 X0))))))\Rightarrow (r1_xxreal_0 np_1 (k15_glib_000 X0))$$