

t25\_nat\_1

(TMFv4VX7ftv9QH4A29E72wRwQSUG8a3g2HR)

October 27, 2020

Let  $v7\_ordinal1 : \iota \Rightarrow o$  be given. Let  $r1\_xxreal\_0 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $np\_1 : \iota$  be given. Let  $k6\_numbers : \iota$  be given. Let  $k1\_nat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $m2\_subset\_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_numbers : \iota$  be given. Let  $k5\_numbers : \iota$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k4\_ordinal1 : \iota$  be given. Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow (\neg (\neg r1\_xxreal\_0 (k1\_nat\_1 X1 np\_1) X0) \wedge ((r1\_xxreal\_0 X1 X0) \wedge (X0 \neq X1)))) \quad (1)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow ((\neg r1\_xxreal\_0 np\_1 X0) \Rightarrow (X0 = k6\_numbers)) \quad (2)$$

Assume the following.

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\forall X1.(v7\_ordinal1 X1) \Rightarrow ((\neg r1\_xxreal\_0 (k1\_nat\_1 X1 np\_1) X0) \Leftrightarrow (r1\_xxreal\_0 X0 X1))) \quad (3)$$

Assume the following.

$$((v2\_xxreal\_0 np\_1) \wedge (m2\_subset\_1 np\_1 k1\_numbers k5\_numbers)) \wedge ((m1\_subset\_1 np\_1 k5\_numbers) \wedge (m1\_subset\_1 np\_1 k1\_numbers)) \quad (4)$$

Assume the following.

$$k5\_numbers = k4\_ordinal1 \quad (5)$$

Assume the following.

$$\forall X0.(m1\_subset\_1 X0 k4\_ordinal1) \Rightarrow (v7\_ordinal1 X0) \quad (6)$$

**Theorem 1**

$$\forall X0.(v7\_ordinal1 X0) \Rightarrow (\neg (r1\_xxreal\_0 X0 np\_1) \wedge ((X0 \neq k6\_numbers) \wedge (X0 \neq np\_1)))$$