

t25_nfcont_3

(TMYfVRbG8hhxM9FAvgJvGmSHaqHzbVz3MVa)

October 27, 2020

Let $v2_struct_0 : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_normsp_0 : \iota \Rightarrow o$ be given. Let $v4_normsp_0 : \iota \Rightarrow o$ be given. Let $v2_normsp_1 : \iota \Rightarrow o$ be given. Let $l1_normsp_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_numbers : \iota$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_rcomp_1 : \iota \Rightarrow o$ be given. Let $v1_nfcont_3 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_nfcont_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k7_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k5_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k7_relat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. (v1_relat_1 X1) \Rightarrow (k9_xtuple_0 (k5_relat_1 X1 X0) = k3_xboole_0 (k9_xtuple_0 X1) X0) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Rightarrow (k3_xboole_0 X0 X1 = X0) \quad (2)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge ((v13_algstr_0 X0) \wedge ((v2_rlvect_1 X0) \wedge ((v3_rlvect_1 X0) \wedge ((v4_rlvect_1 X0) \wedge ((v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge ((v7_rlvect_1 X0) \wedge ((v8_rlvect_1 X0) \wedge ((v3_normsp_0 X0) \wedge ((v4_normsp_0 X0) \wedge ((v2_normsp_1 X0) \wedge (l1_normsp_1 X0)))))))))) \Rightarrow \\ & (\forall X1. ((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k1_numbers (u1_struct_0 X0)))))) \Rightarrow (((v1_rcomp_1 (k1_relset_1 k1_numbers X1) \wedge (v1_nfcont_3 (k2_partfun1 k1_numbers (u1_struct_0 X0) X1 (k1_relset_1 k1_numbers X1)) X0)) \Rightarrow (v1_nfcont_1 (k2_relset_1 (u1_struct_0 X0) X1) X0))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.(v1_relat_1 X1)\Rightarrow(k10_xtuple_0 (k5_relat_1 X1 X0) = k7_relat_1 X1 X0) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(k7_relset_1 X0 X1 X2 X3 = k7_relat_1 X2 X3) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v5_relat_1 X1 X0))\Rightarrow(k2_relset_1 X0 X1 = k10_xtuple_0 X1) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow(k2_partfun1 X0 X1 X2 X3 = k5_relat_1 X2 X3) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1 X1)\wedge(v4_relat_1 X1 X0))\Rightarrow(k1_relset_1 X0 X1 = k9_xtuple_0 X1) \quad (8)$$

Assume the following.

$$\forall X0.(v1_relat_1 X0)\Rightarrow(k5_relat_1 X0 (k9_xtuple_0 X0) = X0) \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((v1_funct_1 X2)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))\Rightarrow((v1_funct_1 (k2_partfun1 X0 X1 X2 X3))\wedge(m1_subset_1 (k2_partfun1 X0 X1 X2 X3) (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.k3_xboole_0 X0 X1 = k3_xboole_0 X1 X0 \quad (11)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow((v4_relat_1 X2 X0)\wedge(v5_relat_1 X2 X1)) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (13)$$

Theorem 1

$$\begin{aligned} & \forall X0. ((\neg v2_struct_0 X0) \wedge (v13_algstr_0 X0) \wedge (v2_rlvect_1 \\ & X0) \wedge (v3_rlvect_1 X0) \wedge (v4_rlvect_1 X0) \wedge (v5_rlvect_1 X0) \wedge \\ & ((v6_rlvect_1 X0) \wedge (v7_rlvect_1 X0) \wedge (v8_rlvect_1 X0) \wedge (v3_normsp_0 \\ & X0) \wedge (v4_normsp_0 X0) \wedge (v2_normsp_1 X0) \wedge (l1_normsp_1 X0)))))) \Rightarrow \\ & (\forall X1. ((v1_funct_1 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k1_numbers (u1_struct_0 X0)))))) \Rightarrow (\forall X2. (m1_subset_1 X2 \\ & (k1_zfmisc_1 k1_numbers)) \Rightarrow (((r1_tarski X2 (k1_relset_1 k1_numbers \\ & X1)) \wedge (v1_rcomp_1 X2) \wedge (v1_nfcont_3 (k2_partfun1 k1_numbers \\ & (u1_struct_0 X0) X1 X2) X0))) \Rightarrow (v1_nfcont_1 (k7_relset_1 k1_numbers \\ & (u1_struct_0 X0) X1 X2) X0)))) \end{aligned}$$