

t25_pepin
(TMK52aRXp7LBTjn9vivsgxQyngdFcwyyveHA)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $np_1 : \iota$ be given. Let $k6_numbers : \iota$ be given. Let $k1_newton : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\neg(k6_numbers \neq X0) \wedge (r1_xxreal_0 X0 k6_numbers)) \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (r1_xxreal_0 k6_numbers X0) \quad (2)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\neg(r1_xxreal_0 X0 np_1) \wedge ((X0 \neq k6_numbers) \wedge (X0 \neq np_1))) \quad (3)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((\neg r1_xxreal_0 np_1 X0) \Rightarrow (X0 = k6_numbers)) \quad (4)$$

Assume the following.

$$\forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (((r1_xxreal_0 np_1 X0) \wedge (r1_xxreal_0 np_1 X1)) \Rightarrow (r1_xxreal_0 X0 (k1_newton X0 X1)))) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1 X0) \wedge (v7_ordinal1 X1)) \Rightarrow (v7_ordinal1 (k1_newton X0 X1)) \quad (6)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v1_xreal_0 X0) \quad (7)$$

Theorem 1

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(v7_ordinal1 X1) \Rightarrow (\neg (\neg r1_xxreal_0 X0 np_1) \wedge ((\neg r1_xxreal_0 X1 k6_numbers) \wedge (r1_xxreal_0 (k1_newton X0 X1) np_1))))$$