

t25_relset_1

(TMYvs2oADwQBZPPJ1SS14vMGx7mD32T14gv)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_relset_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v5_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k10_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $v4_relat_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.(k4_tarski\ X0\ X1 \in k2_zfmisc_1\ X2\ X3) \Leftrightarrow ((X0 \in X2) \wedge (X1 \in X3)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.(m1_subset_1\ X0\ (k1_zfmisc_1\ X1)) \Leftrightarrow (r1_tarski\ X0\ X1) \quad (2)$$

Assume the following.

$$\forall X0.\forall X1.(X0 \in X1) \Rightarrow (m1_subset_1\ X0\ X1) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1_relat_1\ X1) \wedge (v5_relat_1\ X1\ X0)) \Rightarrow (k2_relset_1\ X0\ X1 = k10_xtuple_0\ X1) \quad (4)$$

Assume the following.

$$\forall X0.\forall X1.(r1_tarski\ X0\ X1) \Leftrightarrow (\forall X2.(X2 \in X0) \Rightarrow (X2 \in X1)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.(X1 = k10_xtuple_0\ X0) \Leftrightarrow (\forall X2.(X2 \in X1) \Leftrightarrow (\exists X3.k4_tarski\ X3\ X2 \in X0)) \quad (6)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1\ X2\ (k1_zfmisc_1\ (k2_zfmisc_1\ X0\ X1))) \Rightarrow ((v4_relat_1\ X2\ X0) \wedge (v5_relat_1\ X2\ X1)) \quad (7)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1)))\Rightarrow(v1_relat_1 X2) \quad (8)$$

Theorem 1

$$\begin{aligned} & \forall X0.(\neg v1_xboole_0 X0)\Rightarrow(\forall X1.(\neg v1_xboole_0 X1)\Rightarrow \\ & (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X1 X0)))\Rightarrow \\ & (\forall X3.(X3 \in k2_relset_1 X0 X2)\Leftrightarrow(\exists X4.(m1_subset_1 \\ & X4 X1)\wedge(k4_tarski X4 X3 \in X2)))))) \end{aligned}$$