

t25_toprns_1
(TMG5aMdejnsSApDX6nuk28k4CKM7rqGDL3s7)

October 27, 2020

Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k12_euclid : \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k2_finseq_1 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_euclid : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v4_funct_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.((v1_relat_1 X0) \wedge ((v1_funct_1 X0) \wedge ((v1_finseq_1 X0) \wedge (v3_valued_0 X0)))) \Rightarrow (r1_xxreal_0 k6_numbers (k12_euclid X0)) \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid X0))) \Rightarrow ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 (k2_finseq_1 X0) k1_numbers) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 (k2_finseq_1 X0) k1_numbers)))))) \quad (2)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (u1_struct_0 (k15_euclid X0) = k1_euclid X0) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (4)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (5)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (6)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (7)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (v4_funct_1 (u1_struct_0 (k15_euclid X0))) \quad (8)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (9)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (10)$$

Assume the following.

$$\forall X0.(v4_funct_1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 X0) \Rightarrow (v1_relat_1 X1) \wedge (v1_funct_1 X1)) \quad (11)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid X0))) \Rightarrow (v3_valued_0 X1)) \quad (12)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid X0))) \Rightarrow (v1_finseq_1 X1)) \quad (13)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (14)$$

Theorem 1

$$\forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid X0))) \Rightarrow (r1_xxreal_0 k6_numbers (k12_euclid X1)))$$