

t25_xtuple_0 (TMVzCs-
bGPP7z8oW1ZT5ruwQjr1VsqWhJxMc)

October 27, 2020

Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_xboole_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_xtuple_0 : \iota \Rightarrow \iota$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0. \forall X1. \forall X2. (X2 = k4_xboole_0 X0 X1) \Leftrightarrow (\forall X3. \\ (X3 \in X2) \Leftrightarrow ((X3 \in X0) \wedge (\neg X3 \in X1))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. k4_tarski X0 X1 = k2_tarski (k2_tarski X0 X1) (k1_tarski X0) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (r1_tarski X0 X1) \Leftrightarrow (\forall X2. (X2 \in X0) \Rightarrow (X2 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0. \forall X1. (X1 = k9_xtuple_0 X0) \Leftrightarrow (\forall X2. (X2 \in X1) \Leftrightarrow (\exists X3. k4_tarski X2 X3 \in X0)) \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. k2_tarski X0 X1 = k2_tarski X1 X0 \quad (5)$$

Theorem 1

$$\begin{aligned} \forall X0. \forall X1. r1_tarski (k4_xboole_0 (k9_xtuple_0 X0) \\ (k9_xtuple_0 X1)) (k9_xtuple_0 (k4_xboole_0 X0 X1)) \end{aligned}$$