

t25\_xxreal\_3 (TM-  
MJN8GBLfw6NvVtkBRz6L7y2Ae2Q5L2hh3)

October 27, 2020

Let  $v1\_xxreal\_0 : \iota \Rightarrow o$  be given. Let  $k2\_xxreal\_3 : \iota \Rightarrow \iota$  be given. Let  $k1\_xxreal\_3 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k3\_xxreal\_3 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (k2\_xxreal\_3 (k1\_xxreal\_3 X0 X1) = k1\_xxreal\_3 (k2\_xxreal\_3 X0) (k2\_xxreal\_3 X1))) \quad (1)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (v1\_xxreal\_0 (k2\_xxreal\_3 X0)) \quad (2)$$

Assume the following.

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (k3\_xxreal\_3 X0 X1 = k1\_xxreal\_3 X0 (k2\_xxreal\_3 X1))) \quad (3)$$

Assume the following.

$$\forall X0.\forall X1.((v1\_xxreal\_0 X0) \wedge (v1\_xxreal\_0 X1)) \Rightarrow (k1\_xxreal\_3 X0 X1 = k1\_xxreal\_3 X1 X0) \quad (4)$$

**Theorem 1**

$$\forall X0.(v1\_xxreal\_0 X0) \Rightarrow (\forall X1.(v1\_xxreal\_0 X1) \Rightarrow (k2\_xxreal\_3 (k1\_xxreal\_3 X0 X1) = k3\_xxreal\_3 (k2\_xxreal\_3 X1) X0))$$