

t26_euclid_2

(TMMmurernSVi7S8aHF5h2x4A1iPSD8F9dqD)

October 27, 2020

Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $k23_rvsum_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_rlvect_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k9_binop_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_monoid_0 : \iota \Rightarrow o$ be given. Let $v5_rltopsp1 : \iota \Rightarrow o$ be given. Let $v2_pre_topc : \iota \Rightarrow o$ be given. Let $v13_algstr_0 : \iota \Rightarrow o$ be given. Let $v2_rlvect_1 : \iota \Rightarrow o$ be given. Let $v3_rlvect_1 : \iota \Rightarrow o$ be given. Let $v4_rlvect_1 : \iota \Rightarrow o$ be given. Let $v5_rlvect_1 : \iota \Rightarrow o$ be given. Let $v6_rlvect_1 : \iota \Rightarrow o$ be given. Let $v7_rlvect_1 : \iota \Rightarrow o$ be given. Let $v8_rlvect_1 : \iota \Rightarrow o$ be given. Let $l2_algstr_0 : \iota \Rightarrow o$ be given. Let $l2_struct_0 : \iota \Rightarrow o$ be given. Let $l1_algstr_0 : \iota \Rightarrow o$ be given. Let $l1_rlvect_1 : \iota \Rightarrow o$ be given. Let $l1_rltopsp1 : \iota \Rightarrow o$ be given. Let $l1_pre_topc : \iota \Rightarrow o$ be given. Let $l1_struct_0 : \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v3_valued_0 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $v1_monoid_0 : \iota \Rightarrow o$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1 X0) \Rightarrow & (\forall X1.(m1_subset_1 X1 (u1_struct_0 \\ & (k15_euclid X0))) \Rightarrow (\forall X2.(m1_subset_1 X2 (u1_struct_0 (\\ & k15_euclid X0))) \Rightarrow (\forall X3.(m1_subset_1 X3 (u1_struct_0 (k15_euclid \\ X0))) \Rightarrow & (k23_rvsum_1 (k3_rlvect_1 (k15_euclid X0) X1 X2) X3 = k9_binop_2 \\ & (k23_rvsum_1 X1 X3) (k23_rvsum_1 X2 X3)))))) \end{aligned} \quad (1)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((v2_monoid_0 (k15_euclid X0)) \wedge (v5_rltopsp1 (k15_euclid X0))) \quad (2)$$

Assume the following.

$$\begin{aligned} \forall X0.(v7_ordinal1 X0) \Rightarrow & ((v2_pre_topc (k15_euclid X0)) \wedge \\ & ((v13_algstr_0 (k15_euclid X0)) \wedge ((v2_rlvect_1 (k15_euclid X0)) \wedge \\ & ((v3_rlvect_1 (k15_euclid X0)) \wedge ((v4_rlvect_1 (k15_euclid X0)) \wedge \\ & ((v5_rlvect_1 (k15_euclid X0)) \wedge ((v6_rlvect_1 (k15_euclid X0)) \wedge \\ & ((v7_rlvect_1 (k15_euclid X0)) \wedge ((v8_rlvect_1 (k15_euclid X0)) \wedge \\ & (v5_rltopsp1 (k15_euclid X0)))))))))) \end{aligned} \quad (3)$$

Assume the following.

$$\forall X0.(l2_algstr_0 X0) \Rightarrow ((l2_struct_0 X0) \wedge (l1_algstr_0 X0)) \quad (4)$$

Assume the following.

$$\forall X0.(l1_rlvect_1 X0) \Rightarrow (l2_algstr_0 X0) \quad (5)$$

Assume the following.

$$\forall X0.(l1_rltopsp1 X0) \Rightarrow ((l1_rlvect_1 X0) \wedge (l1_pre_topc X0)) \quad (6)$$

Assume the following.

$$\forall X0.(l1_pre_topc X0) \Rightarrow (l1_struct_0 X0) \quad (7)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.\forall X2.(((v2_rlvect_1 X0) \wedge (l1_algstr_0 \\ & X0)) \wedge ((m1_subset_1 X1 (u1_struct_0 X0)) \wedge (m1_subset_1 X2 (u1_struct_0 \\ & X0)))) \Rightarrow (m1_subset_1 (k3_rlvect_1 X0 X1 X2) (u1_struct_0 X0)) \end{aligned} \quad (8)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow ((v5_rltopsp1 (k15_euclid X0)) \wedge (l1_rltopsp1 (k15_euclid X0))) \quad (9)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.(((v1_relat_1 X0) \wedge (v1_funct_1 X0) \wedge ((\\ & v3_valued_0 X0) \wedge (v1_finseq_1 X0)))) \wedge ((v1_relat_1 X1) \wedge ((v1_funct_1 \\ & X1) \wedge ((v3_valued_0 X1) \wedge (v1_finseq_1 X1)))) \Rightarrow (k23_rvsum_1 X0 \\ & X1 = k23_rvsum_1 X1 X0) \end{aligned} \quad (10)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid X0))) \Rightarrow (v3_valued_0 X1)) \quad (11)$$

Assume the following.

$$\forall X0.(l1_struct_0 X0) \Rightarrow ((v2_monoid_0 X0) \Rightarrow (v1_monoid_0 X0)) \quad (12)$$

Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid X0))) \Rightarrow (v1_finseq_1 X1)) \quad (13)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_monoid_0 X0) \wedge (l1_struct_0 X0)) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 X0)) \Rightarrow ((v1_relat_1 X1) \wedge (v1_funct_1 \\ & X1))) \end{aligned} \quad (14)$$

Theorem 1

$$\begin{aligned} \forall X0.(v7_ordinal1\ X0) \Rightarrow (\forall X1.(m1_subset_1\ X1\ (u1_struct_0 \\ (k15_euclid\ X0))) \Rightarrow (\forall X2.(m1_subset_1\ X2\ (u1_struct_0\ (\\ k15_euclid\ X0))) \Rightarrow (\forall X3.(m1_subset_1\ X3\ (u1_struct_0\ (k15_euclid \\ X0))) \Rightarrow (k23_rvsum_1\ X1\ (k3_rlvect_1\ (k15_euclid\ X0)\ X2\ X3) = k9_binop_2 \\ (k23_rvsum_1\ X1\ X2)\ (k23_rvsum_1\ X1\ X3)))))) \end{aligned}$$