

## t26\_funct\_2

(TMTvqkq2z8qUa2TfJGYC7XCBCVudMjaj6ck)

October 27, 2020

Let  $v1\_funct\_1 : \iota \Rightarrow o$  be given. Let  $v1\_funct\_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $m1\_subset\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_zfmisc\_1 : \iota \Rightarrow \iota$  be given. Let  $k2\_zfmisc\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v2\_funct\_1 : \iota \Rightarrow o$  be given. Let  $k1\_xboole\_0 : \iota$  be given. Let  $k1\_funct\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $k2\_funct\_1 : \iota \Rightarrow \iota$  be given. Let  $v1\_relat\_1 : \iota \Rightarrow o$  be given. Let  $k9\_xtuple\_0 : \iota \Rightarrow \iota$  be given. Let  $k3\_relat\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v4\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Let  $k1\_relset\_1 : \iota \Rightarrow \iota \Rightarrow \iota$  be given. Let  $v5\_relat\_1 : \iota \Rightarrow \iota \Rightarrow o$  be given. Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v1\_funct\_1 X1)) \Rightarrow (((v2\_funct\_1 X1) \wedge (X0 \in k9\_xtuple\_0 X1)) \Rightarrow ((X0 = k1\_funct\_1 (k2\_funct\_1 X1) (k1\_funct\_1 X1 X0)) \wedge (X0 = k1\_funct\_1 (k3\_relat\_1 X1 (k2\_funct\_1 X1) X0)))$$
(1)

Assume the following.

$$\forall X0. \forall X1. ((v1\_relat\_1 X1) \wedge (v4\_relat\_1 X1 X0)) \Rightarrow (k1\_relset\_1 X0 X1 = k9\_xtuple\_0 X1)$$
(2)

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (((X1 \neq k1\_xboole\_0) \Rightarrow ((v1\_funct\_2 X2 X0 X1) \Leftrightarrow (X0 = k1\_relset\_1 X0 X2))) \wedge ((X1 = k1\_xboole\_0) \Rightarrow ((v1\_funct\_2 X2 X0 X1) \Leftrightarrow (X2 = k1\_xboole\_0))))$$
(3)

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow ((v4\_relat\_1 X2 X0) \wedge (v5\_relat\_1 X2 X1))$$
(4)

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1\_subset\_1 X2 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))) \Rightarrow (v1\_relat\_1 X2)$$
(5)

**Theorem 1**

$$\forall X0.\forall X1.\forall X2.\forall X3.((v1\_funct\_1 X3)\wedge ((v1\_funct\_2 X3 X0 X1)\wedge(m1\_subset\_1 X3 (k1\_zfmisc\_1 (k2\_zfmisc\_1 X0 X1))))))\Rightarrow(((v2\_funct\_1 X3)\wedge(X2 \in X0))\Rightarrow((X1 = k1\_xboole\_0)\vee(k1\_funct\_1 (k2\_funct\_1 X3) (k1\_funct\_1 X3 X2) = X2)))$$