

t26_partit_2 (TMKCB-
HVsZyLNN73mHbZaPafYMZaEyS3qy1)

October 27, 2020

Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v8_relat_2 : \iota \Rightarrow o$ be given. Let $r8_relat_2 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k4_tarski : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\begin{aligned} \forall X0.(v1_relat_1 X0) \Rightarrow ((v8_relat_2 X0) \Leftrightarrow (\forall X1.\forall X2. \\ \forall X3.((k4_tarski X1 X2 \in X0) \wedge (k4_tarski X2 X3 \in X0)) \Rightarrow (k4_tarski \\ X1 X3 \in X0))) \end{aligned} \tag{1}$$

Assume the following.

$$\begin{aligned} \forall X0.(v1_relat_1 X0) \Rightarrow (\forall X1.(r8_relat_2 X0 X1) \Leftrightarrow (\forall X2. \\ \forall X3.\forall X4.((X2 \in X1) \wedge ((X3 \in X1) \wedge ((X4 \in X1) \wedge ((k4_tarski \\ X2 X3 \in X0) \wedge (k4_tarski X3 X4 \in X0)))))) \Rightarrow (k4_tarski X2 X4 \in X0))) \end{aligned} \tag{2}$$

Assume the following.

$$\begin{aligned} \forall X0.\forall X1.\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 \\ (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \end{aligned} \tag{3}$$

Theorem 1

$$\begin{aligned} \forall X0.(\neg v1_xboole_0 X0) \Rightarrow (\forall X1.(\neg v1_xboole_0 X1) \Rightarrow \\ (\forall X2.(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X0))) \Rightarrow \\ ((v8_relat_2 X2) \Rightarrow (r8_relat_2 X2 X1)))) \end{aligned}$$