

t26_setlim_1 (TMHDg-
PxQXWgxxw9KRyvYtfgF2P4s6jEjnVvP)

October 27, 2020

Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k5_numbers : \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k9_setfam_1 : \iota \Rightarrow \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $r1_tarski : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_prob_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k3_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_setlim_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_numbers : \iota$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_setlim_1 : \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v3_prob_1 : \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg (X0 \in X1) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 X2)) \wedge (v1_xboole_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (2)$$

Assume the following.

$$\forall X0. (v7_ordinal1 X0) \Rightarrow (r1_xxreal_0 k6_numbers X0) \quad (3)$$

Assume the following.

$$m1_subset_1 k1_xboole_0 k4_ordinal1 \quad (4)$$

Assume the following.

$$\forall X0. \forall X1. (((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k9_setfam_1 X0)))))) \Rightarrow (k3_funct_2 k5_numbers (k9_setfam_1 X0) (k2_setlim_1 X0 X1) k6_numbers = k3_prob_1 X0 X1)) \quad (5)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0)\wedge((\neg v1_xboole_0 X1)\wedge(m1_subset_1 X1 (k1_zfmisc_1 X0))))\Rightarrow(\forall X2.(m2_subset_1 X2 X0 X1)\Leftrightarrow(m1_subset_1 X2 X1)) \quad (6)$$

Assume the following.

$$\forall X0.k9_setfam_1 X0 = k1_zfmisc_1 X0 \quad (7)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (8)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (9)$$

Assume the following.

$$\forall X0.\forall X1.\forall X2.\forall X3.((\neg v1_xboole_0 X0)\wedge(((v1_funct_1 X2)\wedge((v1_funct_2 X2 X0 X1)\wedge(m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))))))\wedge(m1_subset_1 X3 X0)))\Rightarrow(k3_funct_2 X0 X1 X2 X3 = k1_funct_1 X2 X3) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers (k9_setfam_1 X0))\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k9_setfam_1 X0))))))\Rightarrow(k2_setlim_1 X0 X1 = k1_setlim_1 X1) \quad (11)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1)\wedge(v3_ordinal1 k4_ordinal1) \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.((v1_funct_1 X1)\wedge((v1_funct_2 X1 k5_numbers (k9_setfam_1 X0))\wedge(m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k9_setfam_1 X0))))))\Rightarrow((v1_funct_1 (k1_setlim_1 X1)\wedge((v1_funct_2 (k1_setlim_1 X1) k5_numbers (k9_setfam_1 X0))\wedge(v3_prob_1 (k1_setlim_1 X1)))) \quad (13)$$

Assume the following.

$$m2_subset_1 k6_numbers k1_numbers k5_numbers \quad (14)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (15)$$

Assume the following.

$$\begin{aligned} & \forall X0. \forall X1. ((v1_funct_1 X1) \wedge ((v1_funct_2 X1 k5_numbers \\ & (k9_setfam_1 X0)) \wedge (m1_subset_1 X1 (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0)))))) \Rightarrow ((v1_funct_1 (k2_setlim_1 \\ & X0 X1)) \wedge ((v1_funct_2 (k2_setlim_1 X0 X1) k5_numbers (k9_setfam_1 \\ & X0)) \wedge (m1_subset_1 (k2_setlim_1 X0 X1) (k1_zfmisc_1 (k2_zfmisc_1 \\ & k5_numbers (k9_setfam_1 X0)))))) \end{aligned} \quad (16)$$

Assume the following.

$$\forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v1_relat_1 (k1_setlim_1 X0)) \wedge (v1_funct_1 (k1_setlim_1 X0))) \quad (17)$$

Assume the following.

$$\begin{aligned} & \forall X0. ((v1_relat_1 X0) \wedge (v1_funct_1 X0)) \Rightarrow ((v3_prob_1 X0) \Leftrightarrow \\ & (\forall X1. (m2_subset_1 X1 k1_numbers k5_numbers) \Rightarrow (\forall X2. \\ & (m2_subset_1 X2 k1_numbers k5_numbers) \Rightarrow ((r1_xxreal_0 X1 X2) \Rightarrow \\ & (r1_tarski (k1_funct_1 X0 X1) (k1_funct_1 X0 X2)))))) \end{aligned} \quad (18)$$

Assume the following.

$$\forall X0. (m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (19)$$

Assume the following.

$$\forall X0. \forall X1. \forall X2. (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 X0 X1))) \Rightarrow (v1_relat_1 X2) \quad (20)$$

Theorem 1

$$\begin{aligned} & \forall X0. (m1_subset_1 X0 k5_numbers) \Rightarrow (\forall X1. \forall X2. \\ & ((v1_funct_1 X2) \wedge ((v1_funct_2 X2 k5_numbers (k9_setfam_1 X1)) \wedge \\ & (m1_subset_1 X2 (k1_zfmisc_1 (k2_zfmisc_1 k5_numbers (k9_setfam_1 \\ & X1)))))) \Rightarrow (r1_tarski (k3_prob_1 X1 X2) (k3_funct_2 k5_numbers \\ & (k9_setfam_1 X1) (k2_setlim_1 X1 X2) X0))) \end{aligned}$$