

t26_sppol_2
(TMZH328M7SpAXGFYbrWymznbA3iyXtYkj9T)

October 27, 2020

Let $m2_finseq_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $np_2 : \iota$ be given. Let $r1_xxreal_0 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k3_finseq_1 : \iota \Rightarrow \iota$ be given. Let $v2_topreal1 : \iota \Rightarrow o$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $v1_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xreal_0 : \iota \Rightarrow o$ be given. Let $v2_xxreal_0 : \iota \Rightarrow o$ be given. Let $k2_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $np_0 : \iota$ be given. Let $k4_xcmplx_0 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $k1_nat_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_relat_1 : \iota \Rightarrow o$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Let $v3_membered : \iota \Rightarrow o$ be given. Let $v3_xxreal_0 : \iota \Rightarrow o$ be given. Let $v1_xcmplx_0 : \iota \Rightarrow o$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_finseq_1 : \iota \Rightarrow o$ be given. Let $k9_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_topreal1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k1_tarski : \iota \Rightarrow \iota$ be given. Let $k7_partfun1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v2_membered : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0. \forall X1. \forall X2. \neg (X0 \in X1) \wedge ((m1_subset_1 X1 (k1_zfmisc_1 X2)) \wedge (v1_xboole_0 X2)) \quad (1)$$

Assume the following.

$$\forall X0. (v1_xxreal_0 X0) \Rightarrow (\forall X1. (v1_xxreal_0 X1) \Rightarrow (\forall X2. (v1_xxreal_0 X2) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (r1_xxreal_0 X1 X2)) \Rightarrow (r1_xxreal_0 X0 X2)))) \quad (2)$$

Assume the following.

$$\forall X0. \forall X1. (m1_subset_1 X0 X1) \Rightarrow ((v1_xboole_0 X1) \vee (X0 \in X1)) \quad (3)$$

Assume the following.

$$\forall X0. (v1_xreal_0 X0) \Rightarrow (\forall X1. (v1_xreal_0 X1) \Rightarrow (((r1_xxreal_0 X0 X1) \wedge (v2_xxreal_0 X0)) \Rightarrow (v2_xxreal_0 X1))) \quad (4)$$

Assume the following.

$$\begin{aligned} & \forall X0.(v1_xreal_0 X0) \Rightarrow (\forall X1.(v1_xreal_0 X1) \Rightarrow (\forall X2. \\ & (v1_xreal_0 X2) \Rightarrow ((r1_xxreal_0 (k2_xcmplx_0 X0 X1) X2) \Leftrightarrow (r1_xxreal_0 \\ & X0 (k6_xcmplx_0 X2 X1)))))) \end{aligned} \quad (5)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_2) \wedge (m2_subset_1 np_2 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_2 k5_numbers) \wedge (m1_subset_1 np_2 k1_numbers)) \end{aligned} \quad (6)$$

Assume the following.

$$\begin{aligned} & ((v2_xxreal_0 np_1) \wedge (m2_subset_1 np_1 k1_numbers k5_numbers)) \wedge \\ & ((m1_subset_1 np_1 k5_numbers) \wedge (m1_subset_1 np_1 k1_numbers)) \end{aligned} \quad (7)$$

Assume the following.

$$\begin{aligned} & (m2_subset_1 np_0 k1_numbers k5_numbers) \wedge ((m1_subset_1 np_0 \\ & k5_numbers) \wedge (m1_subset_1 np_0 k1_numbers)) \end{aligned} \quad (8)$$

Assume the following.

$$k4_xcmplx_0 np_0 = np_0 \quad (9)$$

Assume the following.

$$k6_xcmplx_0 np_2 np_2 = np_0 \quad (10)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (11)$$

Assume the following.

$$\begin{aligned} & \forall X0.\forall X1.((v7_ordinal1 X0) \wedge (m1_subset_1 X1 k5_numbers)) \Rightarrow \\ & (k1_nat_1 X0 X1 = k2_xcmplx_0 X0 X1) \end{aligned} \quad (12)$$

Assume the following.

$$\forall X0.\forall X1.v1_relat_1 (k2_zfmisc_1 X0 X1) \quad (13)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (14)$$

Assume the following.

$$v3_membered k1_numbers \quad (15)$$

Assume the following.

$$\forall X0.\forall X1.((v7_ordinal1\ X0)\wedge(v7_ordinal1\ X1))\Rightarrow(v7_ordinal1\ (k2_xcmplx_0\ X0\ X1)) \quad (16)$$

Assume the following.

$$\forall X0.((\neg v3_xxreal_0\ X0)\wedge(v1_xreal_0\ X0))\Rightarrow((v1_xcmplx_0\ (k4_xcmplx_0\ X0))\wedge(\neg v2_xxreal_0\ (k4_xcmplx_0\ X0))) \quad (17)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0\ X0)\wedge((\neg v1_xboole_0\ X1)\wedge(m1_subset_1\ X1\ (k1_zfmisc_1\ X0))))\Rightarrow(\forall X2.(m2_subset_1\ X2\ X0\ X1)\Rightarrow(m1_subset_1\ X2\ X0)) \quad (18)$$

Assume the following.

$$\forall X0.\forall X1.(m2_finseq_1\ X1\ X0)\Rightarrow((v1_funct_1\ X1)\wedge((v1_finseq_1\ X1)\wedge(m1_subset_1\ X1\ (k1_zfmisc_1\ (k2_zfmisc_1\ k5_numbers\ X0))))) \quad (19)$$

Assume the following.

$$m1_subset_1\ k5_numbers\ (k1_zfmisc_1\ k1_numbers) \quad (20)$$

Assume the following.

$$\forall X0.((v1_relat_1\ X0)\wedge((v1_funct_1\ X0)\wedge(v1_finseq_1\ X0)))\Rightarrow(m2_subset_1\ (k3_finseq_1\ X0)\ k1_numbers\ k5_numbers) \quad (21)$$

Assume the following.

$$\forall X0.(m2_finseq_1\ X0\ (u1_struct_0\ (k15_euclid\ np_2)))\Rightarrow((v2_topreal1\ X0)\Leftrightarrow(\forall X1.(v7_ordinal1\ X1)\Rightarrow(((r1_xxreal_0\ np_1\ X1)\wedge(r1_xxreal_0\ (k1_nat_1\ X1\ np_2)\ (k3_finseq_1\ X0)))\Rightarrow(k9_subset_1\ (u1_struct_0\ (k15_euclid\ np_2))\ (k2_topreal1\ np_2\ X0\ X1)\ (k2_topreal1\ np_2\ X0\ (k1_nat_1\ X1\ np_1)) = k1_tarski\ (k7_partfun1\ (u1_struct_0\ (k15_euclid\ np_2))\ X0\ (k1_nat_1\ X1\ np_1)))))) \quad (22)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k4_ordinal1)\Rightarrow(v7_ordinal1\ X0) \quad (23)$$

Assume the following.

$$\forall X0.(v1_xreal_0\ X0)\Rightarrow(v1_xxreal_0\ X0) \quad (24)$$

Assume the following.

$$\forall X0.(v3_membered\ X0)\Rightarrow(v2_membered\ X0) \quad (25)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(v1_xxreal_0\ X0) \quad (26)$$

Assume the following.

$$\forall X0.(v7_ordinal1\ X0)\Rightarrow(v1_xreal_0\ X0) \quad (27)$$

Assume the following.

$$\forall X0.(v1_relat_1\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ (k1_zfmisc_1\ X0))\Rightarrow(v1_relat_1\ X1)) \quad (28)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k5_numbers)\Rightarrow(\neg v3_xxreal_0\ X0) \quad (29)$$

Assume the following.

$$\forall X0.(m1_subset_1\ X0\ k1_numbers)\Rightarrow(v1_xreal_0\ X0) \quad (30)$$

Assume the following.

$$\forall X0.(v2_membered\ X0)\Rightarrow(\forall X1.(m1_subset_1\ X1\ X0)\Rightarrow(v1_xxreal_0\ X1)) \quad (31)$$

Theorem 1

$$\forall X0.(m2_finseq_1\ X0\ (u1_struct_0\ (k15_euclid\ np_2)))\Rightarrow((r1_xxreal_0\ (k3_finseq_1\ X0\ np_2))\Rightarrow(v2_topreal1\ X0))$$