

t26_toprns_1 (TMUvThyago-
moj4PxYRsPX1X7x4Y9XL4h4aw)

October 27, 2020

Let $m2_subset_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k1_numbers : \iota$ be given. Let $k5_numbers : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $u1_struct_0 : \iota \Rightarrow \iota$ be given. Let $k15_euclid : \iota \Rightarrow \iota$ be given. Let $k12_euclid : \iota \Rightarrow \iota$ be given. Let $k4_algstr_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v7_ordinal1 : \iota \Rightarrow o$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k4_ordinal1 : \iota$ be given. Let $v3_ordinal1 : \iota \Rightarrow o$ be given. Assume the following.

$$\forall X0.(v7_ordinal1 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (u1_struct_0 (k15_euclid X0))) \Rightarrow (k12_euclid (k4_algstr_0 (k15_euclid X0) X1) = k12_euclid X1)) \quad (1)$$

Assume the following.

$$\forall X0.\forall X1.((\neg v1_xboole_0 X0) \wedge ((\neg v1_xboole_0 X1) \wedge (m1_subset_1 X1 (k1_zfmisc_1 X0)))) \Rightarrow (\forall X2.(m2_subset_1 X2 X0 X1) \Leftrightarrow (m1_subset_1 X2 X1)) \quad (2)$$

Assume the following.

$$k5_numbers = k4_ordinal1 \quad (3)$$

Assume the following.

$$(\neg v1_xboole_0 k4_ordinal1) \wedge (v3_ordinal1 k4_ordinal1) \quad (4)$$

Assume the following.

$$m1_subset_1 k5_numbers (k1_zfmisc_1 k1_numbers) \quad (5)$$

Assume the following.

$$\forall X0.(m1_subset_1 X0 k4_ordinal1) \Rightarrow (v7_ordinal1 X0) \quad (6)$$

Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (\forall X1.(m1_subset_1 X1 (k1_zfmisc_1 X0)) \Rightarrow (v1_xboole_0 X1)) \quad (7)$$

Theorem 1

$$\begin{aligned} & \forall X0.(m2_subset_1 X0 k1_numbers k5_numbers) \Rightarrow (\forall X1. \\ & (m1_subset_1 X1 (u1_struct_0 (k15_euclid X0))) \Rightarrow (k12_euclid (\\ & k4_algstr_0 (k15_euclid X0) X1) = k12_euclid X1)) \end{aligned}$$