

t26_twoscomp
(TML31HACdPQzH2R23EaFC95DihVMqR27PJV)

October 27, 2020

Let $k1_funct_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k33_twoscomp : \iota$ be given. Let $k11_finseq_1 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_numbers : \iota$ be given. Let $np_1 : \iota$ be given. Let $v1_xboole_0 : \iota \Rightarrow o$ be given. Let $k1_xboole_0 : \iota$ be given. Let $k2_binarith : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k8_margrel1 : \iota$ be given. Let $k7_margrel1 : \iota$ be given. Let $np_0 : \iota$ be given. Let $k3_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_xcmplx_0 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k2_xboolean : \iota$ be given. Let $k1_xboolean : \iota$ be given. Let $m1_subset_1 : \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k6_margrel1 : \iota$ be given. Let $k10_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_xboolean : \iota \Rightarrow o$ be given. Let $k3_xboolean : \iota \Rightarrow \iota$ be given. Let $k4_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $v1_funct_1 : \iota \Rightarrow o$ be given. Let $v1_funct_2 : \iota \Rightarrow \iota \Rightarrow \iota \Rightarrow o$ be given. Let $k4_finseq_2 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $np_3 : \iota$ be given. Let $k1_zfmisc_1 : \iota \Rightarrow \iota$ be given. Let $k2_zfmisc_1 : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k7_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k6_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Let $k5_xboolean : \iota \Rightarrow \iota \Rightarrow \iota$ be given. Assume the following.

$$\forall X0.(v1_xboole_0 X0) \Rightarrow (X0 = k1_xboole_0) \quad (1)$$

Assume the following.

$$k2_binarith k8_margrel1 k7_margrel1 = k8_margrel1 \quad (2)$$

Assume the following.

$$v1_xboole_0 np_0 \quad (3)$$

Assume the following.

$$k3_xcmplx_0 np_1 np_0 = np_0 \quad (4)$$

Assume the following.

$$k6_xcmplx_0 np_1 np_1 = np_0 \quad (5)$$

Assume the following.

$$k8_margrel1 = k2_xboolean \quad (6)$$

Assume the following.

$$k7_margrel1 = k1_xboolean \quad (7)$$

Assume the following.

$$k6_numbers = k1_xboole_0 \quad (8)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1 X0 k6_margrel1)\wedge(m1_subset_1 X1 k6_margrel1))\Rightarrow(k2_binarith X0 X1 = k10_xboolean X0 X1) \quad (9)$$

Assume the following.

$$\forall X0.(v1_xboolean X0)\Rightarrow(k3_xboolean (k3_xboolean X0) = X0) \quad (10)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean X0)\wedge(v1_xboolean X1))\Rightarrow(k4_xboolean X0 X0 = X0) \quad (11)$$

Assume the following.

$$v1_xboolean k2_xboolean \quad (12)$$

Assume the following.

$$v1_xboolean k1_xboolean \quad (13)$$

Assume the following.

$$m1_subset_1 k8_margrel1 k6_margrel1 \quad (14)$$

Assume the following.

$$m1_subset_1 k7_margrel1 k6_margrel1 \quad (15)$$

Assume the following.

$$(v1_funct_1 k33_twoscomp)\wedge((v1_funct_2 k33_twoscomp (k4_finseq_2 np_3 k6_margrel1) k6_margrel1)\wedge(m1_subset_1 k33_twoscomp (k1_zfmisc_1 (k2_zfmisc_1 (k4_finseq_2 np_3 k6_margrel1) k6_margrel1)))) \quad (16)$$

Assume the following.

$$\forall X0.(v1_xboolean X0)\Rightarrow(\forall X1.(v1_xboolean X1)\Rightarrow(k7_xboolean X0 X1 = k4_xboolean (k6_xboolean X0 X1) (k6_xboolean X1 X0))) \quad (17)$$

Assume the following.

$$\forall X0.(v1_xboolean X0)\Rightarrow(\forall X1.(v1_xboolean X1)\Rightarrow(k6_xboolean X0 X1 = k5_xboolean (k3_xboolean X0) X1)) \quad (18)$$

Assume the following.

$$\forall X0.(v1_xboolean\ X0) \Rightarrow (\forall X1.(v1_xboolean\ X1) \Rightarrow (k5_xboolean\ X0\ X1 = k3_xboolean\ (k4_xboolean\ (k3_xboolean\ X0)\ (k3_xboolean\ X1)))) \quad (19)$$

Assume the following.

$$\forall X0.(v1_xboolean\ X0) \Rightarrow (\forall X1.(v1_xboolean\ X1) \Rightarrow (k4_xboolean\ X0\ X1 = k3_xcmplx_0\ X0\ X1)) \quad (20)$$

Assume the following.

$$\forall X0.(v1_xboolean\ X0) \Rightarrow (k3_xboolean\ X0 = k6_xcmplx_0\ np_1\ X0) \quad (21)$$

Assume the following.

$$\begin{aligned} & \forall X0.((v1_funct_1\ X0) \wedge ((v1_funct_2\ X0\ (k4_finseq_2\ np_3 \\ & k6_margrel1)\ k6_margrel1) \wedge (m1_subset_1\ X0\ (k1_zfmisc_1\ (k2_zfmisc_1 \\ & (k4_finseq_2\ np_3\ k6_margrel1)\ k6_margrel1)))) \Rightarrow ((X0 = k33_twoscomp) \Leftrightarrow \\ & (\forall X1.(m1_subset_1\ X1\ k6_margrel1) \Rightarrow (\forall X2.(m1_subset_1 \\ & X2\ k6_margrel1) \Rightarrow (\forall X3.(m1_subset_1\ X3\ k6_margrel1) \Rightarrow (k1_funct_1 \\ & X0\ (k11_finseq_1\ X1\ X2\ X3) = k2_binarith\ (k2_binarith\ X1\ X2)\ X3)))))) \quad (22) \end{aligned}$$

Assume the following.

$$k2_xboolean = np_1 \quad (23)$$

Assume the following.

$$k1_xboolean = k6_numbers \quad (24)$$

Assume the following.

$$\forall X0.(v1_xboolean\ X0) \Rightarrow (\forall X1.(v1_xboolean\ X1) \Rightarrow (k10_xboolean\ X0\ X1 = k3_xboolean\ (k7_xboolean\ X0\ X1))) \quad (25)$$

Assume the following.

$$\forall X0.\forall X1.((v1_xboolean\ X0) \wedge (v1_xboolean\ X1)) \Rightarrow (k5_xboolean\ X0\ X1 = k5_xboolean\ X1\ X0) \quad (26)$$

Assume the following.

$$\forall X0.\forall X1.((m1_subset_1\ X0\ k6_margrel1) \wedge (m1_subset_1\ X1\ k6_margrel1)) \Rightarrow (k2_binarith\ X0\ X1 = k2_binarith\ X1\ X0) \quad (27)$$

Theorem 1

$$\begin{aligned} & (k1_funct_1\ k33_twoscomp\ (k11_finseq_1\ k6_numbers\ k6_numbers \\ & k6_numbers) = k6_numbers) \wedge ((k1_funct_1\ k33_twoscomp\ (k11_finseq_1 \\ & k6_numbers\ k6_numbers\ np_1) = np_1) \wedge ((k1_funct_1\ k33_twoscomp \\ & (k11_finseq_1\ k6_numbers\ np_1\ k6_numbers) = np_1) \wedge ((k1_funct_1 \\ & k33_twoscomp\ (k11_finseq_1\ k6_numbers\ np_1\ np_1) = k6_numbers) \wedge \\ & ((k1_funct_1\ k33_twoscomp\ (k11_finseq_1\ np_1\ k6_numbers\ k6_numbers) = \\ & np_1) \wedge ((k1_funct_1\ k33_twoscomp\ (k11_finseq_1\ np_1\ k6_numbers \\ & np_1) = k6_numbers) \wedge ((k1_funct_1\ k33_twoscomp\ (k11_finseq_1 \\ & np_1\ np_1\ k6_numbers) = k6_numbers) \wedge (k1_funct_1\ k33_twoscomp \\ & (k11_finseq_1\ np_1\ np_1\ np_1) = np_1)))))) \end{aligned}$$